

# The Southeastern Bering Sea: Recent Advances in Knowledge

## A Synthesis of Contributions from the

### Southeastern Bering Sea Carrying Capacity Program

### Bering Sea FOCI

and

### The Inner Front Program

### The SEBSCC Synthesis Working Group

George L. Hunt, Jr. Chair

#### Participants:

Chris Baier	Nick Bond
Ric Brodeur	Troy Buckley
Lorenzo Ciannelli	Liz Conners
Chuck Fowler	Susan Henrichs
Jerry Hoff	Anne Hollowed
Nancy Kachel	Allen Macklin
Jeff Napp	Jim Overland
Sigrid Salo	Robert Schabetsberger
Jim Schumacher	Beth Sinclair
Alan Springer	Phyllis Stabeno
Al Tyler	Lucy Vlietstra
Terry Whitledge	

## **Executive Summary**

# Introduction

## Brief History and Goals of Recent Research Programs

The Bering Sea is a semi-enclosed sea that connects the North Pacific and Arctic Oceans. The 500 km-wide eastern continental shelf encompasses about one half of its area and supports extraordinarily rich marine resources. These are of vital importance to the economic survival, subsistence, and cultural foundations of the many indigenous people of western Alaska (IARPC, 2001). Marine resources of the Bering Sea include fisheries equal to about one half of the United States' fishery production, about 80% of the seabirds found in the nation's waters, and substantial populations of marine mammals (NRC, 1996; IARPC, 2001). Its fishery landings include walleye pollock (*Theragra chalcogramma*, a nodal species in the shelf ecosystem), salmon, halibut and crab, and generate over 2 billion dollars in revenue each year (IARPC, 2001). It is thus vital to the economic and social well being of the region that we understand the factors that determine the productivity of the Bering Sea. Such information will facilitate the wise exploitation and stewardship of this most important marine ecosystem.

Since the mid-1990s, two coordinated research programs, the Southeast Bering Sea Carrying Capacity (SEBSCC) Project (1996 – 2002), supported by the Coastal Ocean Program of NOAA, and the Inner Front study (1997-2000), supported by the National Science Foundation (Prolonged Production and Trophic Transfer to Predators: Processes at the Inner Front of the S.E. Bering Sea), were active in the southeastern Bering Sea (Macklin et al., 2002). These two studies were complementary, with one focused on the middle and outer shelf (SEBSCC) and the other (Inner Front) on the inner shelf. Collaboration between investigators in the two programs was strong. In addition, the Bering Sea FOCI (Fisheries-Oceanography Coordinated Investigations) program, funded by NOAA's Coastal Ocean Program, was active from 1991 through

1997, and many of the FOCI investigators were also participants in the Inner Front and SEBSCC programs.

The goals of the Bering Sea FOCI program were to understand the factors that control the abundance of fish populations, and in particular, the abundance and stock structure of walleye pollock in the Bering Sea (Macklin, 1999). A specific objective was to reduce uncertainty in managing these fish. The goals of the SEBSCC program were to increase understanding of the southeastern Bering Sea ecosystem, to document the ecological role of juvenile walleye pollock and factors that affect their survival, and to develop and test annual indices of pre-recruit (age-1) pollock abundance (Macklin et al., 2002). SEBSCC focused on four central scientific issues: 1) How does climate variability influence the marine ecosystem of the Bering Sea? 2) What limits the growth of fish populations on the eastern Bering Sea shelf? 3) How do oceanographic conditions on the shelf influence distributions of fish and other species? 4) What determines the timing, amount, and fate of primary and secondary production? Underlying these broad goals was a narrower focus on walleye pollock. Of particular concern was the understanding of ecological factors that affect year-class strength and the ability to predict the potential of a year-class at the earliest possible time. The Inner Front program focused on the role of the structural front between the well-mixed waters of the **coastal domain** and the two-layer system of the **middle domain**. Of particular interest was the potential for prolonged post-spring-bloom production at the front and its role in supporting upper trophic level organisms such as juvenile pollock and seabirds. Of concern to both programs was the role of interannual and longer-term variability in marine climates and their effects on the function of sub-arctic marine ecosystems and their ability to support upper trophic level organisms (see also Francis et al., 1998).

### **Structure of Synthesis Report**

In this report, we provide an overview of the contributions of the SEBSCC and Inner Front programs to our understanding of processes in the southeastern Bering Sea. We begin with a brief description of the physical environment and marine climate

of the eastern Bering Sea. We follow with an examination of changes and mechanisms of change for the physical and biological components of the eastern Bering Sea ecosystem. We then discuss several conceptual hypotheses that provide possible avenues toward understanding changes in the amount and fate of production in the southeastern Bering Sea, and how climate change may affect the function of this ecosystem. . Finally, we identify a series of questions that recent work suggests will be important to answer in our quest for a fuller understanding of the function of the eastern Bering Sea in a changing climate.

### **Brief Description of the southeastern Bering Sea**

The Bering Sea consists of a deep central basin, a northwestern shelf in the **Gulf of Anadyr** that reaches south along the **Kamchatka Peninsula**, and a broad eastern shelf that stretches from the **Alaska Peninsular** to Russia and the **Bering Strait** (**Figure 1**). For the purposes of this report, the waters of the eastern Bering Sea can be divided into an Oceanic Regime that occupies the basin and a Shelf Regime that occupies the eastern shelf. The eastern shelf can be further sub-divided into the southeastern shelf and a northeastern shelf, with the dividing line running east-west just south of **St. Matthew Island** from the coast to the shelf edge.

The following overview of the Oceanic Regime is based on the description of Schumacher et al. (In Press). The Oceanic Regime of the eastern basin is influenced by **Alaska Stream Water** that enters the Bering Sea through **Amchitka and Amukta** passes, and turns right to form the Aleutian North Slope Current (ANSC; Reed and Stabeno, 1999) (**Figure 2**). This current in turn provides the major source of water for the **Bering Slope Current** (BSC) that varies between following the depth contours of the eastern shelf northwestward with a regular flow, and becoming an ill-defined, variable flow characterized by numerous eddies and meanders (Stabeno et al., 1999a). The eddies occur not only in water seaward of the eastern shelf (Schumacher and Reed, 1992), but also in waters as shallow as 100-122 m deep (Reed, 1998). These eddies are potentially important as habitat for larval and juvenile pollock, and can carry

these fish, as well as nutrient salts, from the Oceanic Domain into the **Outer Shelf Domain** (Schumacher and Stabeno, 1994, Stabeno et al., 1999a).

The broad continental shelf (up to 500 km wide) of the southeastern Bering Sea is differentiated into three bathymetrically-fixed domains, which include the **Coastal Domain** that extends from the shore to about the 50m isobath, the **Middle Shelf Domain**, between the 50 m and 100m isobaths, and the **Outer Shelf Domain** which ranges from 100 m to 200 m in depth (**Figure 3**) (Iverson et al., 1979b; Coachman, 1986; Schumacher and Stabeno, 1998; Stabeno et al., 2001). The domains are separated by fronts or transition zones, with the narrow (5 to 30 km) **Inner Front** or Structural Front between the Coastal Domain and the Middle Shelf Domain, the wide (> 50 km) middle transition zone between the Middle Shelf Domain and the Outer Shelf Domain, and the **Outer Front** between the Outer Shelf Domain and the waters of the slope. In summer, the Coastal Domain is well mixed to weakly stratified, the Middle Shelf Domain is strongly stratified, and the Outer Shelf Domain has well mixed upper and lower layers with a zone of gradually increasing density between (Schumacher et al., In Press). During summer in the Middle Shelf Domain, the temperature difference between the upper and lower layers can be greater than 8° C, and changes in density are dominated by temperature rather than salinity (Hunt et al., 2002a). In the northeastern Bering Sea, changes in tidal energy and freshwater discharge from the **Yukon River** affect the location of the fronts, with the Inner Front occurring in water 30 m or less (REF ??? Better than Bering Sea Research Plan, page 11). During summer in Norton Sound, a two-layered system can occur in water as shallow as 20 m. In winter, the water column there is well mixed.

### **Climate and the Bering Sea Marine Ecosystem**

An important development in fisheries oceanography during the last decades of the twentieth century was the realization that climate variability, at the scale of decades,

could have profound impacts on the function of marine ecosystems. Of particular importance was the realization that seemingly small shifts in the mean values of atmospheric variables, at least when compared to their interannual variability, could result in major changes in the productivity or standing stocks of fish populations. Recent work has suggested that climate variations may affect the mechanisms (e.g. bottom-up or top-down) that control populations, such that the impact on fish stocks of a given rate of harvest may be quite different in different climate patterns.

### **Climate Indices**

For the North Pacific Ocean, among many potential indices of climate variability, there are three well-studied indices of climate patterns that depend on variability in winter, the Pacific Decadal Oscillation (PDO), the El Niño/Southern Oscillation (ENSO), and the Arctic Oscillation (AO). The PDO is defined as the leading mode of sea surface temperature variability in the North Pacific (north of 20° N), and has time scales of 20 – 30 years (Ladd et al., ms) (**Figure 4**). The PDO is a major mode of variability (Wallace et al., 1992), yet it explains only 21% of the variance of the monthly SST and is primarily centered on the central North Pacific rather than the Gulf of Alaska and Bering Sea. The ENSO has widespread influence on global climate variability at time scales of 2 – 7 years, and has its greatest influence in the tropics. Recently, ENSO has been shown to have a small but significant influence on the marine climate of the Bering Sea via atmospheric teleconnections (Niebauer, 1998; Hollowed et al., 2001; Martin et al., 2001; Overland et al., 2001). The AO is defined as the leading mode of sea level pressure variability north of 20° N, and consists of a pattern of zonally symmetric variability in the strength of the polar vortex (Thompson and Wallace, 1998; Ladd et al., ms). The AO has its largest variance in winter (January – March). The strongest mode of variability in the AO is interannual, but it also varies at decadal scales, having changed sign in 1976 and again in 1989 (Thompson and Wallace, 1998; Overland et al., 1999) (**Figure 4**). The AO has an influence on the Bering Sea through its affect on the Aleutian Low, which is the monthly or seasonal mean location of the center of low sea level pressure over the North Pacific (Overland et al., 1999). The value and position of the Aleutian

Low reflects the strength and distribution of storm tracks in the southern Bering Sea and sub-arctic Pacific Ocean. These storms have great influence on the marine climate of the Bering Sea in winter.

A fourth index of atmospheric pressure, the North Pacific (NP) pattern, represents the leading mode in spring of the 700 mb height and is most prominent from March through July (Barnston and Livezey, 1987; Ladd et al., ms). The NP consists of a north-south pressure dipole, and its strong variance in spring relates to storminess in the Bering Sea (Overland et al., 2002).

In the North Pacific region, the wintertime indices, in conjunction with indices of biological responses in marine ecosystems, have been used to identify abrupt shifts in climatic forcing and ecosystem response at decadal time scales (e.g., Trenberth and Hurrell, 1995; Mantua et al., 1997; Francis et al., 1998; Hare and Mantua, 2000; McFarlane et al., 2000; Hollowed et al., 2001). Two of these regime shifts have been identified in the past thirty years. One followed the winter of 1976-1977, in which the PDO and the AO both shifted (**Figure 4**). A second shift, of just the AO, occurred after the winter of 1988-1989 (Ebbesmeyer et al., 1991; Hare and Francis, 1995; Sugimoto and Tadokoro, 1998; Beamish et al., 1999; Brodeur et al., 1999a; Hare and Mantua, 2000). There is some evidence of a third shift in the winter of 1998-1999 (Schwing and Moore, 2000; Peterson et al., 2002). Although the ENSO appears to alternate between two states that are repeatedly visited, that does not appear to be the case for regime shifts in the southeastern Bering Sea, where the few regimes documented so far have each had unique characteristics.

The main climate feature influencing the southeastern Bering Sea is the Aleutian Low (**Figure 5**). The Bering Sea lies between the cold Arctic air mass to the north and warmer maritime air mass of the North Pacific. Over the Bering Sea, there is a region of strong gradients in sea-level pressure (SLP) between the Aleutian Low and the high pressure over the Arctic. Considerable interest has developed around low frequency

(multi-decadal) variability in the North Pacific in both the physical and biological portions of the North Pacific ecosystem (Minobe, 1999). Hare and Mantua (2000) found evidence in 100 time-series for regime-like jumps in these records near 1977 and 1989. The evidence was clearer in the biological data than in the physical data. Their second mode showed strong covariability between physical and biological variations in the Bering Sea over the past 40 years.

To investigate this variability, the strength of the Aleutian Low is plotted in **Figure 6** for 1900 through 2000. Visually, there are some indications of a lowering of SLP after 1927, higher pressures from 1947 – 1977, and an event near 1990. However, the main feature of this plot is the large change from year to year. In fact, almost two-thirds of the variance in this record is at time scales shorter than five years (Overland et al., 1999). When one looks at the time series after 1977, almost half the years have SLP values near or slightly above the mean (horizontal line) that occur between years of record low SLP. Using a five-year running mean is not particularly representative of individual years, either for periods when the values were near average or for those periods with large negative anomalies as the averaged data remove the signal from the extreme years that may force the system.

One approach to modeling the variability of the Aleutian low time series is to fit its auto-correlation function, the correlation of the time series with itself as a function of lag between the series. **Figure 7** shows the auto-correlation function (the vertical stick plots) repeated in all three plots as a function of lag in years. The values are all small, generally less than 0.3; however, the values are mostly positive, suggesting a broad contribution of frequencies to the original time series, rather than a Gaussian distribution of random errors. The solid heavy lines represent three models of the time series (Percival et al., 2001). The left plot is an Auto-Regressive (AR) red noise model that rapidly decays to zero. The middle plot is a long memory (LM) model that rapidly decays in the first years but then has a broad contribution from longer lags. The third model (right) is a 50-year oscillation plus white noise. All three models are fit with two

free parameters and all are candidates to represent the Aleutian Low, because a 100-year record is too short an interval to say one model is superior based on classical statistical grounds.

We have additional information that favors the long memory model. The LM model assumes contributions from many processes with different time lags, while the AR model represents a single time scale. It is more reasonable to expect that atmospheric variability over the Bering Sea is influenced by many processes, such as North Pacific SST, Arctic air masses, ENSO, and Siberian storm systems. We also know from the work of Hare and Mantua (2000) that regime-like behavior favors the LM and oscillator models, as the LM model is about five times more likely to have a 20-year run of all positive or negative values than the AR model.

What does this mean for the biology of the Bering Sea? Instead of a pure regime shift model of the physical system with changes spread over 20+ years, we have a physical system which has a strong response in one or two years. The low frequencies contribute to the timing of these strong events. In this conceptual model, the impact of the change in the physical system is large enough to promote the reorganization of the ecosystem. The massive increase in pollock after 1978 and 1989 (Wespestad et al., 2000) and the 1997 coccolithophore event (Sukhanova and Flint, 1998; Vance et al., 1998; Overland et al., 2001) follow these patterns. Thus, the regimes seen in the biological data of Hare and Mantua (2000) may be the result of a continuing decadal ecosystem reorganization following a major meteorological extreme.

### **Meteorological Forcing**

The variability in SST warming is expected to be a dominant factor on the Bering Sea shelf during the warm season. **Figure 8** summarizes the 40-year record of the rate of warming in early summer and 1 August SST (top panel); two crucial components of the surface heating, the downward shortwave (solar) radiation and latent heat fluxes (middle panel); and two aspects of the wind, the rate of wind mixing ( $u_x^3$ ) (**Figure 8**,

bottom panel), and along-shelfbreak wind stress (**Figure 9, bottom panel**). These time series reveal that recent August SSTs are roughly 1° C warmer than those typifying the 1960s. Much of the trend in this heating can be attributed to a long-term tendency towards more solar heating and decreasing surface latent heat fluxes, or evaporative cooling, over the 40-year record. Interannual variability in the summertime warming appears to be due to a combination of variations in solar heating and wind mixing.

The physical state of the ocean over the shelf during spring and summer has been well documented only since 1995, which compromises interpretation of decadal-scale changes in the marine ecosystem. Recent work has indicated that there exists a viable approach for addressing this lack of direct oceanographic observations for retrospective analyses. Specifically, it appears that reasonably accurate hindcasts of the evolution of thermal profiles over the shelf can be made using a 1-D ocean model (the Price-Weller-Pinkel or PWP model) forced by surface winds and heat fluxes generated from reanalysis of historical records (Ladd et al., ms). These hindcasts can be used to estimate not just temperatures (an important factor for zooplankton growth rates) but also the supply of nutrients to the euphotic zone, with implications for summertime primary production.

Progress has been made in understanding the impact of climate variations on the Bering Sea shelf, but there remain a host of important but unanswered questions. For example, while the mean flow on the shelf is sluggish, it is likely at times to be significant. It remains unclear which aspects of atmospheric forcing are important to cross-shelf flow, and what time scales are of primary importance to these events. In general, the sources and potential predictability of climate variations for the Bering Sea in spring through fall are poorly understood. In particular, little attention has been devoted to air-sea interactions during fall and the possible ramifications of their variability for the ecosystem. Thus, while there is a growing appreciation that relatively short-lived but intense events can account for a disproportionate share of the seasonally

integrated forcing, it remains unknown how the frequency and timing of these events relate to the longer-term variations in climate.

## **Physical Components of the Eastern Bering Sea Ecosystem**

### **Sea Ice in the Eastern Bering Sea**

Sea ice extent is a crucial aspect of the physical environment of the eastern Bering Sea shelf. The Bering Sea is a marginal ice zone, which is typically ice-free from June through October. Beginning in November, cold winds from the Arctic cool the water and begin the formation of ice in the polynyas that form in the lees of islands and coasts. Throughout winter, the prevailing winds advect the ice southward into warmer water where the ice melts at its southern edge, cooling and freshening the seawater (Pease, 1980; Niebauer et al., 1999). Recently, Niebauer (1998) has found that the position and depth (strength) of the Aleutian Low has a significant effect on the sea ice cover of the Bering Sea, and that the effect of the Aleutian Low is linked to ENSO events. The maximum southerly extent of the ice and the amount of ice melt affect fluxes of heat and salt, thereby influencing both baroclinic flow and the temperature at the bottom (the cold pool) in the Middle Shelf Domain (Ohtani and Azumaya, 1995; Schumacher and Stabeno, 1998; Wyllie-Echeverria and Wooster, 1998).

One of the more readily observed impacts of climate change on the eastern Bering Sea ecosystem is the extent and duration of sea ice over the Bering Sea shelf (Niebauer, 1998; Stabeno et al., 2001; Hunt et al., 2002a). The seasonal variation in the position of the ice edge is about 1700 km, the most extensive of any Arctic or sub-arctic region (Niebauer, 1998). In an average winter, about 75% of the eastern shelf is ice covered, but the amount and duration of ice cover can vary interannually by up to 25% of the seasonal range, depending on the wind field (Niebauer, 1983; 1998; Schumacher and Stabeno, 1998). During the last two decades, the maximum ice extent over the eastern shelf occurred, on average, in March, but maximum ice extents have

been as early as January (in 2000), and as late as the end of April (in 1976) (Stabeno and Hunt, 2002). During the early and mid-1970s, ice arrived early over the southeastern shelf and persisted into spring. Following this cold period, there was a warmer period when sea ice was less common. Finally, in 1989 there appeared to be a shift to cooler conditions although not as cold as those observed in 1972–1976 (Stabeno et al., 2001).

During cold winters, ice can cover most of the eastern shelf (Stabeno et al., 2001). Alternately during warmer winters, ice does not extend much farther south than **St. Matthew Island**. Analysis of ice charts (**Figure 10**) shows that decadal patterns of variability are evident (Niebauer, 1998; Hunt et al., 2002a). In the period from 1977 to 1996, there was a 5% reduction of ice cover as compared to 1947-1977 (Niebauer, 1998). Interannual and decadal-scale variability in sea-ice coverage was greater at the southernmost edge of the ice field than farther north (**Figure 10**). Between 57° and 58° N, there was an apparent decrease in days with ice after 1 January between 1972 - 1976 (mean number of days with ice  $130 \pm 18$  SD) as compared with 1977 - 1989 ( $67 \pm 26$  SD,  $t = 1.767$ ,  $p = 0.096$ ) and 1990 - 2000 ( $76 \pm 23.3$  SD,  $t = 2.036$ ,  $p = 0.061$ ). The average pattern of ice coverage has also changed since the early 1970s (**Figure 11**), when the maximum extent of the zone in which  $\geq 10\%$  ice cover was present annually for  $> 2$  weeks extended farther south and west than it did in the 1980s or the 1990s (Stabeno et al., 2001). Additionally, in the 1980s and 1990s, the zone where ice lasted for at least 2 weeks withdrew northeastward along the **Alaska Peninsula**. Most of the north side of the Peninsula has been ice-free since 1990. The 1989 regime shift did not result in a return to the extensive ice conditions present before 1977.

In the eastern Bering Sea, a proxy for sea ice extent is provided by the mean winter (Jan-Apr) surface temperature (Bond and Adams, 2002). The 40-year record for the site of Mooring 2 (top panel of **Figure 9**) shows a notably cold period in the early to middle 1970s, and a warm period in the late 1970s into early 1980s, but conditions during the 1990s are similar to those in the 1960s. The time series of atmospheric

parameters directly related to sea ice (middle panel of **Figure 9**) indicate that the presence of sea ice is a function of not just the cross-shelf component of the wind, but is also strongly related to the net surface heat fluxes. Niebauer (1998) found that before the regime shift of 1976/77, below-normal ice cover in the eastern Bering Sea was associated with El Niño conditions, during which the center of the Aleutian Low shifted eastward, and resulted in warm air from the Pacific flowing over the southeastern Bering Sea. After the regime shift, the Aleutian Low was located even farther eastward during El Niño periods, and under these circumstances the southeastern shelf was subjected to north and east winds from the interior of Alaska, which resulted in increased ice cover.

An examination of the timing of the ice retreat and air temperature at 500 hPa (~ 5 km altitude) reveals that during the last decade there has been a marked change in the timing of spring transitions (Stabeno and Overland, 2001). While sea ice has extended farther south in the last decade, it has retreated more quickly, resulting in the northern Bering Sea being ice free earlier than in previous decades. In addition, atmospheric temperature during May has increased by 3°C in the 1990s compared to the 1980s.

### **Biological Importance of the Timing of Ice Retreat**

In the southeastern Bering Sea, the timing of spring primary production is determined by a combination of the date of ice retreat, stabilization of the water column by solar heating and the cessation of strong storm activity (Sambrotto et al., 1986; Stabeno et al., 1998, 2001; Eslinger and Iverson, 2001). The timing of the spring bloom is important because it determines the ambient water temperatures in which grazers of the bloom must forage. Data illustrative of the conditions that determine the timing of the spring bloom were obtained from Mooring 2, located in ~ 72 m of water in the middle domain (**Figures 1, 12**). If ice retreat comes before mid-March, there is insufficient light to support net production in the well-mixed water column (e.g., **Figure 12**, 1996, 1998, 2000). Without ice remaining after mid-March, the spring bloom is delayed until May or

June, after winter winds have ceased and thermal stratification stabilizes the water column (**Figure 12**) (Stabeno et al., 1998, 2001; Eslinger and Iverson, 2001). If ice melt occurs in April or May, there is an early, ice-associated bloom (e.g., **Figure 12**, 1995, 1997, 1999). The pattern of late ice retreats with early ice-related blooms and early ice retreats with late blooms has held since the 1970s (**Table 1**). Although wind mixing of the water plays a role in determining when the bloom will occur, it is apparent in **Figure 13** that the timing of the last winter storm is less important than the date of ice retreat in determining the timing of the bloom. Thus, early blooms occur in cold water and are related to ice-edge blooms, whereas late blooms occur in relatively warm water and are not related to the ice edge (**Figures 12, 14**).

### **Cold Pool Formation**

The bottom waters over the eastern Bering Sea shelf show considerable interannual variation in temperature. Over the northern parts of the shelf, cold, salty brine is rejected as sea ice forms in polynyas, and this dense water sinks to the bottom. Density flows to the north carry much of this salty bottom water through Bering Strait to the Arctic Ocean where it contributes to the halocline (Cavalieri and Martin, 1994; Schumacher and Stabeno, 1998).

Over the central and southeastern Bering Sea shelf, cold bottom waters are formed when sea ice melts, and the cold, fresh meltwater is mixed throughout the water column by storms (Stabeno et al., 1998). These meltwaters can chill the entire water column to about  $-1.7^{\circ}\text{C}$ . When the surface waters are warmed by solar radiation in spring, a thermocline forms, and the cold bottom waters are largely insulated from further heating (Coachman, et al., 1980; Ohtani and Azumaya, 1995; Wyllie-Echeverria, 1995). Bottom temperatures in this “cold pool” warm slightly over the summer, but may remain below  $2^{\circ}\text{C}$  until storm-induced mixing occurs in fall (Ladd et al., ms). The extent and temperature of the southern cold pool is dependent on the amount of ice melt that occurs, and since ice is constantly melting as it is advected south, the amount of

meltwater generated is a function of the duration of time that ice is present (Pease, 1980; Overland and Pease, 1982).

### **Biological Impacts of Water Temperature**

The temperature of the water column beneath the surface is important for determining the rates of the physiological processes of organisms residing there. Physiological processes vary as a power function of temperature, and phytoplankton growth is less sensitive to water temperature than is zooplankton growth (see also Vidal, 1980; Vidal and Smith, 1986; Townsend et al., 1994). Water temperature exerts a strong influence on the growth rates of zooplankton, and is often thought of as more important than food availability for limiting the growth rates of small-bodied copepods (McLaren, 1963; Corkett and McLaren, 1978; Vidal, 1980; Dagg et al., 1984; Huntley and Lopez, 1992). Thus, in years with warm water, Walsh and McRoy (1986) hypothesized that zooplankton would capture more of the primary production than in cold years, and the greater production of zooplankton in warm years would support the pelagic community, e.g., fish such as pollock. Thus, in the middle domain where interannual environmental variability is greatest, water temperature is likely to play a major role in interannual variation in copepod biomass (Smith and Vidal, 1984, 1986; Napp et al., 2000) (**See section on Zooplankton, below**).

Water temperature also affects the timing of hatching and survival of pollock eggs (Blood, 2002). Modeling of hatching times based on laboratory experiments and temperatures found in the Bering Sea predicted that the hatching periods for pollock eggs could vary as much as 13 days between the warmest and coldest years encountered between 1995 and 1998. A longer incubation period means that the eggs are exposed to predation for a greater period and that there is increased risk of exposure to extreme low temperatures as the eggs in the upper water column can be exposed to temperature variations driven by short-term weather fluctuations. Malformation of pollock embryos has been reported for eggs incubated at  $-1^{\circ}\text{C}$

(Nakatani and Maeda, 1984), a temperature not infrequently encountered in the shelf waters of the Bering Sea shelf in late winter and early spring (Hunt et al., 2002a).

The cold pool has important effects on the distribution and survival of fish. For example, juvenile walleye pollock prefer to avoid waters  $< 2^{\circ}\text{C}$  (Wyllie-Echeverria, 1996). When the southern cold pool is of reduced size, these fish spread out over much of the middle domain in shelf waters not frequented by adult pollock. When the southern cold pool is extensive, the juvenile pollock move toward the warmer waters of the outer domain and shelf edge, where they are subject to increased levels of cannibalism by adult pollock that reside in these outer shelf waters (Ohtani and Azumaya, 1995; Wyllie-Echeverria, 1995, 1996; Wyllie-Echeverria and Wooster, 1998).

### **On-shelf Fluxes and Nutrient Replenishment**

On-shelf fluxes of nutrient salts from the basin are critical for the long-term productivity of the eastern Bering Sea shelf, but the mechanisms responsible for forcing these fluxes are still not well understood. In the northern Bering Sea, it has been suggested that water from the basin and slope crosses the shelf to flow northward through **Anadyr Strait** and **Bering Strait (Figure 3, currents)** (Shuert and Walsh, 1993). Nutrients in this water would then be available to support the extraordinarily high rates of summertime production found in the **Chirikov Basin** and northward through Bering Strait (Springer et al., 1996). This process would also provide a mechanism for the transport of large oceanic copepods onto the northern shelf, where they support immense populations of planktivorous seabirds (Springer and Roseneau, 1985; Springer et al., 1987, 1989; Hunt and Harrison, 1990; Hunt et al., 1990; Russell et al., 2001). However, the exact connections to the Bering Sea basin remain unclear, as observations by Stabeno and Reed (1994) and model results (Overland et al., 1994) suggest that the **Bering Slope Current**, the presumed source of water carrying nutrients and copepods to the **Chirikov Basin**, may turn westward south of  $59^{\circ}\text{N}$ . Rather, the source of the slope water passing through **Anadyr Strait** may be water that has advected onto the outer shelf episodically or through the canyons of the shelf edge

as far south as Bering Canyon; these then flow along the outer shelf with increasing intensity to the north and west, and thence through Anadyr Strait (Stabeno et al., 1999a).

Over the southeastern Bering Sea shelf, nutrient replenishment was initially thought to result from tidally driven diffusion (Coachman, 1986). However, more recent work shows that the coefficients required for tidally-driven diffusion are larger than those present on the shelf (Stabeno et al., 2001). Although mean annual current velocities over the middle shelf at Mooring 2 are weak, currents averaged over shorter periods (e.g., daily) can exceed  $25 \text{ cm s}^{-1}$ . Currents are strongest in near-surface waters (34-month mean,  $1.2 \text{ cm s}^{-1}$  at 15 m), and much weaker at the bottom ( $0.2 \text{ cm s}^{-1}$  at 60 m) (Stabeno et al., 2001). Currents are strongest in winter and weakest in summer. In 1998, these currents were sufficiently strong to advect organisms typical of the oceanic regime into Middle Domain waters adjacent to the Inner Front near Cape Newenham and **Nunivak Island** (Hunt et al., 1999, Coyle and Pinchuk 2002b). It is also hypothesized that the generally seaward movement of sea ice in winter may result in an onshore flow at depth that could contribute to replenishment of nutrients over the southeastern shelf, but this hypothesis has yet to be investigated (Schumacher and Alexander, 1999).

Mechanisms for on-shelf transport of nutrients include eddies that bring slope waters onto the shelf at least as far as the 150-m isobath (Stabeno et al., 1999a; Stabeno and Van Meurs, 1999; Okkonen, 2001; Johnson et al., submitted), and Reed (1998) has observed them in waters between 100 and 120m about 20% of the time (**Figure 3**). However, eddies are rare in water < 100 m deep, and other mechanisms are required to replenish nutrients in the Middle and Inner Domains. Two regions of preferential on-shelf flow are **Bering Canyon**, which is just north of the **Aleutian Islands** near **Unimak Pass**, and the area west of the **Pribilof Islands**, where the shelf break narrows (Stabeno et al., 1999a). There, acceleration of flow over the outer shelf (Coachman, 1986; Schumacher and Stabeno, 1998) results in entrainment of slope

water (Stabeno et al., 1999b). On-shelf flow west of the **Pribilof Islands** can move into the Middle Domain where it is marked by a front to the northeast of **St. Paul Island** (Flint et al., 2002), or it may be entrained around the islands in tidal currents (Stabeno et al., 1999b). Two measures of the atmospheric forcing of the ocean circulation, the wind stress curl and the wind stress along the **Alaskan Peninsula/Aleutian Island** chain, also exhibit substantial variability (bottom panel of **Fig. 9**), but it remains an open question whether the flow over the shelf is sensitive to these effects.

### **Stratification, Mixing and the Vertical Flux of Nutrients**

Shelf waters of the southeastern Bering Sea, although well-mixed during winter by storms, stratify in late spring from solar heating (Eslinger and Iverson, 2001; Ladd et al. submitted). This stratification inhibits vertical flux of nutrients, and once the spring bloom has exhausted them from the upper mixed layer, the lack of nutrients limits new production. These nutrients can be replenished when processes break down the stratification. Sambrotto et al. (1986) identified the importance of summer storms as a mechanism for deepening the pycnocline and stirring nutrients into the upper mixed layer where they could be taken up by plant cells in the presence of light. Analysis of wind speed cubed, a measure of the ability of winds to mix the upper water column shows that summer winds have declined since the early 1980s (**Figure 15**). Ladd et al. identified the importance of winter conditions for determining the strength of the pycnocline, and hence the ease with which it could be eroded by storms. A second pathway for the movement of nutrients from depth to the surface layers is upward mixing in the vicinity of fronts (Iverson et al., 1979a, Sambrotto et al., 1986; Kachel et al., 2002). For example, this upward mixing can result in regions of high productivity on the stratified side of the inner front (Hunt et al., 1996a; Kachel et al., 2002). These mechanisms require the availability of a pool of nutrients at depth in the vicinity of the front. In 1997 after the completion of the spring bloom, a severe storm in late May mixed the water column to depths of 65 m or more and resulted in renewed production. The result was a depletion of nutrients to 60 m or more, and a lack of availability of nutrients at the base of the inner front and at the pycnocline for mixing into the upper

mixed layer (Stockwell et al., 2001). This observation pointed to the importance of episodic events in structuring the ecology of shelf waters.

## **Biological Components of the eastern Bering Sea Ecosystem**

### **Primary Production**

During the past decade, we have made considerable advances in our understanding of factors influencing the timing of the spring bloom (see above, Stabeno et al., 2001, Hunt et al., 2002a; Hunt and Stabeno, 2002). However, we know less about the magnitude of the bloom, and whether the amount of production varies between ice-associated and open water blooms. Work by Alexander and Niebauer (1981) and Niebauer et al. (1981, 1990, 1995) suggests that the ice edge-bloom used to be as great or greater than the open water bloom in the early 1980s. In 1997, which had an ice-associated bloom followed by a second, open-water bloom subsequent to a mixing event in May, primary production, based on nutrient draw-down, was greater than in the early 1980s (Stockwell et al., 2001).

The question of whether there has been a change in the amount of net annual new production is important. Schell (2000), using stable isotope ratios from carbon sequestered in the baleen of bowhead whales during periods of feeding in the northern Bering Sea, estimated that primary production in the Bering Sea had decreased by as much as 30 to 40% since 1967, with almost all of the decrease coming since 1976. Cullen et al. (2001) questioned whether some of this effect was the result of anthropogenic CO<sub>2</sub> or changes in the species composition of the phytoplankton. However, Schell (2001) provided additional information, including data from stable isotopes of nitrogen, that corroborates the earlier findings. Grebmeier and Cooper (1994, 2002), Grebmeier and Dunton (2000), and Grebmeier (1992) have found evidence for declines in sediment oxygen respiration of as high as 73% over the period 1987 to 2002, with declines in benthic biomass of 89% over a longer period. In addition,

they have documented changes in the species composition of benthic bivalves and other fauna. Taken together, these studies point to a decline in production levels for the northern Bering Sea, including in the Saint Lawrence Island Polynya region south of **St. Lawrence Island**. These reductions in production may be related to reduced northward flow through Bering Strait (Roach et al., 1995), and a consequent diminution of nutrient advection from the Bering Sea basin onto the northern shelf.

In the southeastern Bering Sea, there appears to be no clear indication of a decrease in production, though Hirons et al. (2001) have attempted to extend the results of Schell (2000) to the remainder of the eastern Bering Sea and **Gulf of Alaska** by examining stable isotope ratios in the teeth of harbor seals (*Phoca vitulina*), northern fur seals (*Callorhinus ursinus*) and Steller sea lions (*Eumetopias jubatus*). For these species, when data from teeth from both the Gulf of Alaska and the Bering Sea were combined, they found a significant decline in the  $\delta^{13}\text{C}$  in sea lions and similar, though non-significant, trends in harbor seals and fur seals. However, from their published data, it is difficult to tell whether these trends depend upon the combination of data from the Bering Sea and the Gulf of Alaska, or whether the trends would hold up within regions as well.

In contrast, in 1997, estimates of primary production in the southeastern Bering Sea, based on the reduction of nitrate over the middle and inner shelf, suggested that new production may have been between 10 and 30% greater in 1997 than in the early 1980s (Stockwell et al. 2001). Estimates for production levels in 1998 and 1999, however, do not appear to differ from those of the early 1980s (Whitledge, Pers. Comm.). In contradiction to these estimates of productivity in 1997, 1998 and 1999, sediment trap data supported the notion that productivity was higher in 1998 than 1997 (Smith et al., 2002), as did data from the uptake of ammonium (Rho, 2000). However,  $\delta^{13}\text{C}$  values from copepods in the 1997-1999 period were lower than found by Schell et al. (1998). Smith et al. hypothesized that this decline in  $\delta^{13}\text{C}$  could be the result of diminished production, similar to that found by Schell (2000), or it could be the result of

other factors. Interestingly, in the period 1997-1999, Smith et al. did not find a decrease in  $\delta^{13}\text{C}$  in euphausiids and chaetognaths, as would have been expected if the decline in copepod  $\delta^{13}\text{C}$  was the result of changes in primary productivity. Based on the sum of the above results and the high stable or increasing biomass of fish and invertebrates over the southeastern shelf (see below), it seems most unlikely that there has been a marked reduction in primary production there since the 1970s (Hunt et al., 2002a).

### **Coccolithophore Bloom**

In the 1990s, there were marked anomalies in the species composition of phytoplankton in the eastern Bering Sea. Although diatoms typically dominate phytoplankton biomass in the eastern Bering Sea (Sukanova et al., 1999), in the late 1990s, coccolithophore blooms dominated summer phytoplankton assemblages over much of the shelf (**Figures 16, 17**) (Sukhanova and Flint, 1998; Vance et al., 1998; Napp and Hunt, 2001, Stockwell et al., 2001). Coccolithophores are small (5-20  $\mu\text{m}$ ) phytoplankton surrounded by calcium carbonate plates. Coccolithophore blooms are common in the North Sea, Gulf of Maine, and coastal eastern North Pacific, and characteristically occur in nutrient depleted waters with warm, shallow mixed layers (Holligan et al. 1983; Mitchell-Innes & Winter 1987; Balch et al. 1992; Townsend et al. 1994). High densities of coccoliths result in a whitening of the water detectable by satellite imagery (Holligan et al. 1983; Balch et al. 1991; Brown & Yoder 1994; Gower 1997). Although coccolithophore blooms had not been documented previous to 1997 in the eastern Bering Sea, there are satellite images of "white" water from there (Brown & Yoder 1994).

The first coccolithophore bloom (*Emiliana huxleyi*) recorded from the Bering Sea was initially observed 3 July 1997 in the middle domain (Vance et al. 1998), although *E. huxleyi* cells were present in the water as early as May and June (Stockwell et al., 2001). By early August, the bloom was at least 200 km wide (Tynan 1998), and by early September it covered  $2.1 \times 10^5 \text{ km}^2$  of the middle domain and parts of the inner domain (Sukhanova and Flint, 1998; Vance et al., 1998; Napp and Hunt, 2001). It was

apparent in SeaWiFS imagery from 18-25 September and traces could be detected as late as October. This event was unusually large and long-lived when compared with coastal blooms of coccolithophores described previously (**Table 2**) (Holligan et al. 1983; Balch et al. 1991; Brown & Yoder 1994; Gower 1997). Most coastal blooms are on the order of  $10^4$  km<sup>2</sup>, and persist less than 40 days. Concentrations of coccolithophore cells were as great or greater than those reported from the North Atlantic (Townsend et al. 1994; Robertson et al. 1994).

The coccolithophore bloom greatly reduced light penetration and visibility in the water. Near **Nunivak Island (Figure 2)**, the depth of the 1% light level, often taken as the maximum limit for net photosynthesis, shoaled from between 18 and 33 m in June to between 5 and 15 m in September in the bloom (S. Zeeman, Pers. Comm.). At **Slime Bank (Figure 1)**, outside the bloom, the depth of the 1% light level remained the same (27 m). Underwater videos near the **Pribilof Islands** documented cloudy bloom-water ranging in depth from 7 m to 44 m, and extending to the bottom in several locations (R. Brodeur, pers. Comm.).

The eastern Bering Sea coccolithophore bloom recurred yearly from 1997 through 2001, and in most years was comparable in spatial extent and cell density to large-scale coccolithophore blooms in other parts of the world's oceans, though of considerably longer duration (**Table 2**). New algorithms for the analysis of SeaWiFS imagery showed that the coccolithophore blooms started in February as melting began along the ice edge and then spread northward, peaking in April (Iida et al. 2002). The blooms with the largest aerial extent occurred in the warm years of 1998 and 2000, whereas those in the cold years of 1999 and 2001 were smaller (Iida et al., 2002).

The summer of 1997 was marked by unusually warm surface temperatures, a strong thermocline and depletion of nitrate and silicate from the surface waters (Napp and Hunt, 2001). The conditions under which the 1997 bloom commenced were similar to conditions believed to be conducive to coccolithophore blooms elsewhere (Balch et

al. 1992; Holligan et al. 1983 Townsend et al. 1994). However, we do not know why the eastern Bering Sea blooms were initiated, or why they recurred over a several year period when there was great variability in oceanographic conditions (Stabeno and Hunt, 2002). We also lack information on why they ceased to occur in 2002, and on their role in energy flux to the food webs of the eastern Bering Sea shelf.

## **Zooplankton**

Microzooplankton are protists and metazoan organisms smaller than 200 $\mu\text{m}$  that are present in the plankton (Dussart, 1965). Because they are small and individually inconspicuous, their role in the world ocean has been under-appreciated until recently. Although microzooplankton are an abundant element in the food webs of the southeastern Bering Sea, we are only just beginning to investigate their role in this ecosystem (Howell-Kubler et al., 1996; Olson and Strom, 2002). For example, in April, 1992, microprotozoan abundances ranged from 300 to 6233 organisms  $\text{l}^{-1}$  with a biomass of 0.58 to 9.73  $\mu\text{g C l}^{-1}$  (Howell-Kubler et al., 1996). These biomass levels were similar to those of other oceanic regions, and were estimated to be sufficient to meet the metabolic needs of first-feeding larval pollock, though observations to determine if pollock use this resource are not available.

Microzooplankton were an important component of the southeastern Bering Sea food webs in the summer of 1999. Within the coccolithophore bloom, 75 % of the total chlorophyll *a* (Chl *a*) came from cells > 10  $\mu\text{m}$  (mostly the diatom *Nitzschia* spp.), and average growth rates for cells > 10  $\mu\text{m}$  and < 10  $\mu\text{m}$  were nearly equal (Olson and Strom, 2002). Within the bloom, microzooplankton grazing rates were only 28% of the growth rates of phytoplankton growth rates. In contrast, for the shelf as a whole, grazing by microzooplankton accounted for 110% of the growth of cells > 10  $\mu\text{m}$  and only 81% of the growth of cells < 10  $\mu\text{m}$ . This preferential grazing on the larger cells may help to explain the persistence of the coccolithophore bloom and is contrary to the belief that microzooplankton are constrained to diets of nanophytoplankton (Olson and Strom, 2002). These findings emphasize the need for a thorough examination of the

role of microzooplankton both in the summer, for which we have evidence that they may form an important link between phytoplankton and meso-zooplankton, and during the spring bloom, when their role has yet to be evaluated.

Traditionally, calanoid copepods have been believed to be the major agents of energy transfer between large-celled diatoms and upper trophic level consumers such as planktivorous fish (e.g., Hood, 1999). Although emerging data now suggest that in boreal oceans these large copepods have a large dietary component of microzooplankton (Capriulo et al., 1991; Sherr and Sherr, 1992; Rivkin et al., 1999), the large copepods are still important prey for fish, whales and seabirds. In the eastern Bering Sea, the copepod communities of the basin and outer shelf are dominated in spring by large species of *Neocalanus* (*N. cristatus*, *N. plumchrus*, and *N. fleminergi*) and *Eucalanus bungii*, and the middle and inner shelf by the smaller *Calanus marshallae*, *Pseudocalanus* spp. and *Acartia* spp. (Cooney and Coyle, 1982; Smith and Vidal, 1986).

The most complete time series of zooplankton abundances in the eastern Bering Sea depend on data gathered in summer by the T/S *Oshoro Maru*, which has documented declines zooplankton biomass from the basin between the late 1960s and the early 1990s (basin, **Figure 18**) (Sugimoto and Takadoro, 1997). In the basin, they also found a biennial fluctuation in zooplankton biomass that was negatively correlated with the catch of Asian pink salmon (*Oncorhynchus gorbuscha*). Sugimoto and Takadoro interpreted these two patterns as indicating a bottom-up control of the zooplankton on a decadal-scale, and a top-down control on an annual scale. In contrast to the results obtained from the basin, examination of *Oshoro Maru* data from the shelf showed neither a long term trend, nor evidence of biennial cycles in summer zooplankton biomass (**Figure 18**) (Hunt et al., 2002a; Napp et al. 2002). Although there was considerable interannual variation, no discernable temporal trend was detected.

For spring, two sets of studies show that the abundance of small shelf species of copepods varies with sea temperature. In 1980, the upper layer of the middle and outer shelves of the southeastern Bering Sea warmed slowly as compared to 1981 (Smith and Vidal, 1986). In May 1981, small copepods of the middle shelf were more abundant than in 1980 (**Table 3**), and *Calanus marshallae* was observed to have two generations in 1981, rather than the expected one (Smith and Vidal, 1986). In the 1990s, there was also the opportunity to compare a very cold year (1999) with two years in which water temperatures were high (1997, 1998). For species of small copepods over the inner and middle shelf areas, June abundances in 1999 were reduced by up to 90% compared to the two warmer years (**Table 4**) (Coyle and Pinchuk, 2002b). Although Smith and Vidal hypothesized that differences in predation on the copepods as well as temperature might have affected the differences in abundance between 1980 and 1981, there was no indication in the data of Coyle and Pinchuk (2002b) that chaetognaths were responsible for the declines in copepod abundance observed in 1999. Coyle and Pinchuk (2002b) provide compelling evidence that, even on a station-by-station basis, there was a strong relationship between the numbers of copepods present and integrated water temperatures (e.g., **Figure 19**). They estimated that secondary production of calanoids in spring 1999 was about 3-4% that which occurred in the warm years of 1997-1998. Interestingly, by August – September, there were no consistent significant differences in the biomass of small copepods between 1999 and the two warmer years (Coyle and Pinchuk, 2002b).

The effect of temperature on euphausiids appeared to be the inverse of its effect on copepods; in 1999, the acoustically measured biomass of adult euphausiids on the inner and middle shelf was significantly higher than in 1997-1999 (**Table 5**) (Coyle and Pinchuk, 2002a). However, Coyle and Pinchuk point out that this difference may be related to a delay in euphausiid breeding in the cold year, which would result in more adults remaining in the water column when they were sampled in June than in a warm year when most adults would have spawned and died prior to June. Coyle and Pinchuk (2002a) noted that there were significantly higher densities of euphausiid eggs and

larvae present in 1999 (the cold year) compared to the warm years of 1997 and 1998. Coyle and Pinchuk (2002a) concluded that there were no significant differences in the biomass of euphausiids over the inner and middle shelf in 1997 and 1998 compared to earlier periods for which data were available. However, if euphausiid spawning is completed in early spring and there is a reduced availability of late spawning adults in summer, this change in the timing of availability could have a negative impact on predators, such as short-tailed shearwaters (*Puffinus tenuirostris*) that depend on euphausiids for a significant portion of their diet (Baduini et al., 2001a,b; Hunt et al., 2002b).

In the last three decades, gelatinous zooplankton, in particular large scyphomedusae, have gone through a remarkable increase in biomass and then crash (**Figure 20**) (Brodeur et al., 1999a, 2002). The cause (or causes) of the outbreak of jellyfish is not known, though it has been hypothesized that changing climate and ocean temperatures may have been the trigger (Brodeur et al. 1999a). It has also been hypothesized that a decrease in forage fish over the southern portion of the shelf in the early 1980s may have contributed to the jellyfish increase by releasing them from competition (Brodeur et al., 2002). Currently, there are no hypotheses to explain why they suddenly decreased. At the **Pribilof Islands**, the dominant scyphozoan, *Chrysaora melanaster*, was estimated to consume about one third of the standing stock of crustacean zooplankton and 4.7% of their annual production. Additionally, these jellyfish were estimated to consume about 3% of the standing stock of age-0 pollock in the vicinity of the Pribilof Islands (Brodeur et al., 2002).

## **Fish**

During the late-1970s and early 1980s, several stocks of groundfish and non-crab invertebrates showed strong changes in biomass in the eastern Bering Sea (Conners et al., 2002). Conners et al. identified three sites that had been surveyed consistently since about 1965 using comparable gear. Patterns of change in biomass for all species combined were similar across these three areas (Figure 21), and timing

of changes in both commercially exploited and non-exploited species were similar. Pacific cod and several species of flatfish showed changes of 300 to 600 percent, whereas the biomass of Greenland turbot decreased by 90 percent (Figure 22). These changes resulted in a marked shift in the species composition of ground fish and benthic invertebrates in shelf waters. Conners et al. noted that the timing of the change in CPUE in the trawls was consistent with the timing of the major regime shift of 1976/77, and that there was little evidence of responses to the later, weaker, regimes shifts in 1989 and 1998.

The female spawning biomass of the pollock stock increased strongly in the 1980s because of the growth and survival of the strong year classes that started in 1972 (Figure 23). The biomass continued to be above the long-term average during the 1990s because of the strong year-classes that followed the exceptionally strong 1989 year-class. The increases were apparently due to a combination of favorable ocean conditions and the conservative fishery management practices put in place by the North Pacific Fisheries Management Council. The stock showed fluctuations as the young recruits grew in size and were taken by the fishery or died through natural mortality (Figure 22).

Year-class success represents an annual estimate of productivity and survival for a species. During the period from 1963 to 2001, year-class strength of walleye pollock in the eastern Bering Sea varied from a low of 3.6 billion age-1 fish in 1963 to a high of 66.0 billion in 1978 (Figure 23). Hollowed and Wooster (1995) classified years into those warmer and colder than the long term mean for the North Pacific Ocean from 1946 to 1990. They found that during a warm period, the year-class strength of many stocks of groundfish were stronger, while during a cool period, the same stocks showed weaker year class strength. Exceptionally strong year classes of pollock occurred in 10 years (Figure 23, Table 6). There were two or three banner year-classes per decade with intermediate years showing average to weak year classes (NPFMC, 2000).

Other species of groundfish in the eastern Bering Sea showed the same general pattern as walleye pollock of increase in biomass during the early 1980s due to strong year-classes following the 1976/77 climate regime shift. As examples (Figure 22), Pacific cod (*Gadus macrocephalus*), yellowfin sole (*Limanda aspera*), northern rock sole (*Lepidopsetta polyxystra*), Alaska plaice (???) , flathead sole (*Hippoglossoides elassodon*), arrowtooth flounder (*Atheresthes stomias*) and Greenland turbot (*Reinhardtius hippoglossoides*) follow this pattern (NPFMC, 2000). These trends were due to the increased productivity that was manifest as year-class success (Figure 23).

Two alternative recruitment patterns seem to have emerged. There are those species that show strong year classes in the late 1970s and early 1980s, and there are species with a delayed run of strong year classes beginning in the late 1980s and continuing through the early 1990s (Table 6). Both patterns show a reduction of year class strength in the late 1990s. Fishes with strong year-class productivity in the late 1970s and 1980s include Pacific cod, yellowfin sole, flathead sole, and Alaska plaice. Fishes without strong recruitment in the 1970s, but having strong year-classes in the 1980s are rock sole and arrowtooth flounder (Figure 23). Greenland turbot are unique in that they had strong year classes in the late 1970s, but not in the 1980s and 1990s. There are years that stand out as being characterized as having coincident strong or weak year-classes among eight species of groundfish (Table 6). The years in which over half of the species had strong year-classes were 1977 (5/8), 1979 (6/8), and 1984 (5/8). Two years had weak year classes for over half of the species, 1986 (5/8) and 1994 (7/8).

An important difference between walleye pollock and other groundfish is the continued high biomass levels of walleye pollock in the Bering Sea in the 1990s which contrasts with patterns of decline for many other economically important species (Figure 22). Walleye pollock is unique in its continued production of young through the decade of the 1990s (Figure 23, Table 6).

The pattern of changes in biomass available to the fisheries has followed the availability of recruits, not the catch of the fishery. However, these stock changes greatly affected the fishery landings and sustainable yields. The most notable series of stock increases came after the 1976/76 regime shift that resulted in favorable ocean conditions north of the state of Washington. During the 1980s, simulations showed that wind-driven surface currents in the southeastern Bering Sea were mostly eastward, and flatfish as well as pollock enjoyed high levels of recruitment (Wilderbuer et al. (2002). However, in the 1990s, surface currents were westerly (seaward), and flatfish recruitment was weaker. For the flatfish, this would have resulted in advection to favorable nursery grounds (Wilbderbuer et al., 2002); for pollock, advection into Bristol Bay would result in separation of larvae and juveniles from cannibalistic adults (Wespestad et al., 2000).

The late 1970s and early 1980s were also a period of major change in the management of the fisheries. In 1977, the extended fishery jurisdiction of the United States and Canada came into force. International fishing was severely curtailed and fishery management became progressively more conservative. There was a transition period of “joint venture” fishing during the late 1970s and early 1980s, after which only domestic fishing was allowed. Also, because of the cooperative interaction between the International Pacific Halibut Commission and the North Pacific Fisheries Management Council, the catch of cod and flatfish has been conservative. Thus, fishery takes have affected the total biomass of groundfish in the eastern Bering Sea, patterns in change appear to be driven by ecological determinants of recruitment.

Another group of species that are of great importance to both the commercial sector and to subsistence users are the Pacific salmon. The world’s largest runs of wild sockeye salmon (*Oncorhynchus nerka*) occur in Bristol Bay (**REF**), and important runs of other species occur in most of the drainages of western Alaska. Several of the runs showed remarkable increases in the 1970s (e.g., Yukon River chum (*O. keta*) and Bristol Bay sockeye), but returns from these and other stocks in western Alaska plunged

precipitously in the late 1990s (Figure 24). The reasons for these declines are not apparent, but the most likely hypotheses focus on ocean survival (Kruse 1998). In recent years, the ocean ecology of salmon in the eastern Bering Sea has received little attention, and we have few data on their impact on the ecosystem there (e.g., Nishiyama, 1974), nor on factors that might affect their ocean ecology. Work recently begun by the North Pacific Anadromous Fish Commission (the BASIS program), will begin to remedy this lack of information (**REF**).

In addition to work on the commercially important groundfish and salmon species, there have also been significant gains in our knowledge of the distribution, abundance, and ecology of forage fishes. These fish include both small species, such as sand lance (*Ammodytes hexapterus*), capelin (*Mallotus villosus*), Pacific herring (*Clupea pallasii*), smelts, and mesopelagic groups such as Myctophids (lanternfish) and Bathylagidae (deepsea smelts), as well as the young of larger species such as age-0 and age-1 walleye pollock. These species are eaten by large fish, including Pacific cod, Pacific halibut (*Hippoglossus stenolepis*) and walleye pollock, and are also important components of the diets of marine mammals and seabirds. Brodeur et al. (1999b) summarized data collected in 1987 in a joint program with Russian colleagues that resulted in a pair of surveys that covered most of the Bering Sea. They found that age-0 and age-1 pollock were more abundant on the eastern shelf than elsewhere in the Bering, that herring and capelin were most abundant on the northern and western shelves. They also compared the distribution of various species, as sampled in the National Marine Fisheries Bottom Trawl Surveys, in a warm year (1987) and in a cold year (1986). They found that some forage fishes, such as age-1 pollock, avoided regions with cold bottom temperatures, whereas others did not (e.g., capelin).

A group of little studied forage species that deserve particular attention are the mesopelagic fish and squid, which are important forage for upper trophic level consumers including large fish, marine mammals and seabirds. Many mesopelagic species migrate into the upper mixed layer each night to forage on euphausiids and

copepods, returning to depth during the day. This behavior provides a mechanism for rapid downward transport of production from the upper ocean. One species, *Stenobrachius leucopsarus*, perhaps the ecologically most important, may have a biomass of 21 million tones in the North Pacific sub-arctic gyres (including the Bering Sea and the Sea of Okhotsk) (Beamish et al., 1999). Sinclair et al. (1999) assembled data on mesopelagic fishes and squids from the Bering Sea to provide a description of the species composition, distribution and relative abundances of members of this important fauna. They noted that, of the squids, members of the families Gonatidae and Cranchiidae appeared in the highest numbers in both trawl surveys and predator diets, while of the fishes, Myctophidae and Bathylagidae were most common. Sinclair and Stabeno (2002) sampled basin waters near the edge of the southeastern Bering Sea shelf. They found unexpectedly high concentrations of biomass, driven primarily by the deep-sea smelt *Leuroglossus schmidti*. Their work revealed a new species of gonatid, a range extension for the snailfish (*Paraliparis paucidens*), and new size records for several species of fish and squid. Discoveries of this nature demonstrate that this ecologically important group of organisms has been neglected. Given their large biomass compared to that of most fish species, we must devote more effort to learning about them. Because of their strongly diel cycles of behavior, study of their ecological role will require examination of trophic relations both during the daylight and at night.

The waters around the Pribilof Islands have been thought to provide a uniquely productive habitat for juvenile pollock because of the potential for enhance production associated with frontal processes there. During the 1990s, several projects examined the ecology of age-0 walleye pollock there. These are referenced in the section on the Pribilof Islands (Chapter two), and will not be discussed here.

### **Marine mammals and seabirds**

Dramatic population declines of pinnipeds and seabirds in the eastern Bering Sea and Gulf of Alaska since the mid-1970's contrast with their stability elsewhere in the North Pacific Ocean. Previous studies have proposed a combination of natural

(climatic) and anthropogenic (fisheries) factors that may have altered the structure of the food web in the eastern Bering Sea (Piatt and Anderson 1996; NRC 1996; Merrick 1997; Springer 1998; Anderson and Piatt 1999; Hunt et al. 2002a) and thus affected population stability of apex predators. Although the mechanisms of change unique to the Bering Sea are not fully understood (Hunt and Byrd 1999; Hunt et al. 2002a), changes in the populations of marine birds at the Pribilof Islands are likely the result of declines in prey availability, as there is little evidence that these seabird species are subject to significant levels of predation.

#### *Marine mammal Populations –*

Approximately 80% of the world population of northern fur seals (*Callorhinus ursinus*) breeds on the Pribilof Islands where population estimates have been conducted since the turn of the century, and with consistent methodology for the past 60 years (York and Hartley 1981). There are currently an estimated 800,000 animals remaining from an estimated population of 2 million as recently as the 1950's. The species is listed as depleted under the Marine Mammal Protection Act (NMFS 1993). Unlike most other apex predators in the Bering Sea, the pattern of decline among northern fur seals is one of steady to severe reductions in population numbers followed by some years of stability (Figure 25). A steep decline began in the mid-1970's, the effect of either reduced pup production or decreased juvenile survival during their pelagic phase (York and Kozloff 1986). Declines continued into the mid-1980's, after which the St. Paul Island population remained stable at 33% of its 1970's population, while the smaller population of St. George Island stood at 60% of its 1970's level and continued to decline incrementally into the 1990's (York et al. 2000; Loughlin et al., 1994). The most recent population count in 2002 indicates that an annual decline of 5% has been occurring on both islands since 1998 (NMFS 2002).

Steller sea lions (*Eumetopias jubatus*) are currently divided into two stocks. The western stock ranges from 144° W in the Gulf of Alaska westward across the Aleutian Island chain and includes Walrus Island in the Pribilof Island group. After a continual

population decrease of 5% per year since the 1970's, the monitored portion of the stock in the Gulf of Alaska and along the Aleutian Islands chain showed an increase of 5% between 2001 and 2002. The population of Steller sea lions on Walrus Island (Pribilof Island group) has been in steady decline since the 1960's and is considered a nearly extinct rookery (Figure 26). The western stock of Steller sea lions is currently listed as endangered under the U.S. Endangered Species Act. The causes of the decline in western stock of Steller sea lions remains controversial (NRC, 2003).

Populations of large cetaceans in the eastern Bering Sea were greatly reduced by commercial whaling that continued through about 1970 (NRC, 1996). In contrast to declines in pinniped and seabird populations (see below), populations of large baleen whales, which were almost extirpated in shelf waters of the eastern Bering Sea by the 1960s (NRC, 1996; Merrick, 1997), may be increasing. Although in the mid-1970s, it was rare to encounter any large cetacean over the eastern Bering Sea shelf (Leatherwood et al., 1983; Brueggerman et al., 1987), since the 1980s, whales have been encountered over shelf waters with increasing frequency (Baretta and Hunt, 1994). Significant numbers of large cetaceans are now present in shelf waters during summer (Tynan, 1999; Moore et al., 2000, 2002).

#### *Pinniped Diets -*

The composition of the diet of northern fur seals in the Bering Sea has been inferred from material obtained from stomachs of animals shot at sea, from stomach contents and vomits of animals on land, and most recently from analysis of hard parts remaining in scats (feces) (Sinclair et al., In Prep.). Although not completely comparable because of differences in the likelihood of detection of different dietary items, the time series for fur seals provide evidence of significant changes in the composition of their diets over the past forty years (see Hunt et al., 2002a for a review of earlier work). Most significantly, in the 1960s, cold-water species such as capelin and Greenland turbot were important, whereas by the 1980s, these had disappeared from fur seal diets at the Pribilofs and diets were dominated by pollock  $\leq$  2 years of age

During the 1990s, analyses of scats on St. Paul and St George Islands have revealed that fur seals there take at least 4 species of squid and approximately 30 species of fish. Primary prey (those species that occurred with a frequency of 10% or higher on either island, in any single year) included the four species of squid and five species of fish. On St. Paul Island, only walleye pollock ranked above 10% for all sample years combined. In all years combined on St. George Island, pollock, salmon, and the gonatid squid *Gonatopsis borealis/Berryteuthis magister*<sup>1</sup> ranked as primary prey items. Since 1988, the number of northern fur seals on the Pribilof Islands consuming squid has decreased, while the number consuming walleye pollock (*Theragra chalcogramma*) and northern smoothtongue (*Leuroglossus schmidti*) on both islands, and Pacific salmon (*Salmonidae*) on St. George Island has increased (Figure 29). While northern fur seals have relied primarily on juvenile pollock (ages-0 and -1) and to a lesser degree age-2+ since the 1980's (Sinclair et al. 1994; 1996; Antonelis et al. 1997), in the 1990s, there has been an increase in the use of age-3+ pollock, and possibly a decrease in the consumption of age-0 pollock, particularly on St. Paul Island (Figure 30). Over the decade of the 1990s, fur seals on St. George have had a somewhat higher consumption of age-3+ pollock compared to those on St. Paul Is., which is in line with the greater densities of adult pollock around St. George Island relative to St. Paul (NMFS).

Adult fur seals are typically specific to summer feeding locations (Robson 2002) depending upon the rookery of their birth; adult females from St. George Island can be generally characterized as off-shelf feeders compared to those from St. Paul Island (Sinclair et al. 1996; Goebel?). Off-shelf feeding is indicated by consumption of salmon, northern smoothtongue, and the Gonatid squid *Gonatopsis borealis/Berryteuthis magister*.

---

<sup>1</sup> These two species cannot currently be differentiated based on beak structure alone (Sinclair et al. 1994).

Scats collected from Steller sea lions on Walrus Island in 1994 (n=43) and 2001 (n=44) contained 11 species (or taxonomic groups) of fish-prey that ranked above 10% in frequency of occurrence (FO) in one or both years. Irish Lord (*Hemilepidotus hemilepidotus*), Pacific cod, walleye pollock, and rock sole ranked above 10% FO in both sample years. With the exception of salmon, all the species consumed by Steller sea lions on Walrus Island are demersal or semi-demersal in habit. Both juvenile and adult pollock were present in Steller sea lion scats with an emphasis of age 5+ in year two.

#### *Seabird Populations:*

During the 1970s and early 1980s, the abundance of piscivorous seabirds breeding at the Pribilof Islands declined (Figures 27, 28) (Decker et al., 1995, Springer, 1998, Dragoo et al., 2001). Kittiwake (*Rissa* spp.) and common murre (*Uria aalge*) declines were especially severe on St. Paul Island during the 1970s and 1980s. During that time, common murre became more abundant on St. George Island. It is unclear why population declines in murre were more severe on St. Paul Island than on St. George Island.

Schmutz (Appendix 1 in Vlietstra et al. 2003) partitioned the causes of kittiwake and murre population changes on the Pribilof Islands between reproductive success (productivity) and other factors that include post-fledging mortality and emigration. He found that for the four species of kittiwakes and murre, productivity accounted for one-third (34%) of the variation in population change. The majority (66%) of variation in population growth rate was attributable to mortality and emigration, but distinction between these 2 demographic processes was not possible for most species due to lack of data. However, other survival and population data for the geographically restricted Red-legged Kittiwake (*Rissa brevirostris*) indicated that, for this species, variations in survival have likely been a principal demographic force.

#### *Marine bird diets:*

Sampling of marine bird diets has shown that there are consistent, significant differences in diet components between seabirds nesting on St George and St Paul islands, and that, since the mid-1970s, there have been significant changes of diet components for seabird species nesting on St George and St. Paul islands (Vlietstra et al., 2003). During the 1970s and 1980s, black-legged kittiwakes (*R. tridactyla*) and common and thick-billed murres (*U. lomvia*) used more invertebrates (e.g., euphausiids, squid) on St. George than on St. Paul Island. Similarly, during the 1990s, the proportion of black-legged kittiwake and thick-billed murre diets consisting of invertebrates was higher on St. George than on St. Paul Island. During the 1970s and 1980s, there were also differences in the proportions of four species of forage fishes used (Vlietstra et al., 2003). Capelin composed a higher proportion of kittiwake diets on St. Paul Island than on St. George Island. Sand lance were a larger proportion of murre diets on St. George Island than on St. Paul Island. During the 1990s, all four seabird species took a higher percentage of gadids on St. Paul Island than they did on St. George Island.

The differences in diets between the islands are important. They indicate that kittiwakes and murres on St. George have had access to a wider variety of prey, in particular, invertebrates including squids and euphausiids, than seabirds on St. Paul Island. And, seabirds on St. Paul Island have been more dependent on juvenile gadids and capelin than birds on St. George Island where seabirds apparently had more ready access to a wider variety of forage fishes including sand lance and myctophids.

During the 1970s and 1980s, gadids (mainly walleye pollock) were the primary component of black-legged kittiwake, common murre, and thick-billed murre diets. Red-legged kittiwakes took mainly lanternfish. We observed a significant decline between the 1970s and 1980s in the proportion of gadids in kittiwake diets on St Paul, but not St George Island (Figure 31) and in thick-billed murre diets on St. George Island. Likewise, the diet of red-legged kittiwakes during the early-season contained a higher proportion of gadids during the 1970s than they did during the 1980s.

### *Seabird Reproduction in an ecosystem context*

Vlietstra et al. (2003) evaluated the spatial scale at which causal factors appeared to have influenced demographic parameters in Pribilof Islands' seabirds. If population declines were related to processes that operate on large (regional) spatial scales, such as prey availability over the shelf, then seabird reproductive success and colony attendance should have been correlated between St. Paul and St. George Islands. If populations responded to processes that operated on small (local) spatial scales, such as prey availability associated with oceanographic features near an island, then seabird demography should have differed between the islands. Because colony attendance and reproductive success of black-legged kittiwakes and red-legged kittiwakes were correlated between populations on St. Paul and St. George Islands, as were interannual variations in the diets of these species on the two islands, Vlietstra et al. (2003) concluded that population dynamics in kittiwakes were influenced by processes that occurred over large spatial scales (shelf-wide), rather than at the local (island) level. In contrast, for both common and thick-billed murres, interannual variability in colony attendance, and reproductive success in thick-billed murres, were not correlated between islands, suggesting that the sizes of populations of two murres were influenced by factors that operated on a local scale.

At about the time of the 1976/77 regime shift, and concurrent with the declines in piscivorous pinnipeds and seabirds at the Pribilof Islands, were significant decreases in the biomass of age-1 walleye pollock on the Bering Sea shelf (Brodeur et al. 1999b, Hunt et al. 2002a, Vlietstra et al., 2003). In some areas, abundance decreased by as much as 95% (Springer 1992, Sinclair et al. 1996, Brodeur et al. 1999b, Hunt et al., 2002a). Likewise, there is evidence for significant declines in the abundance of capelin, an important forage fish for seabirds (Hunt et al., 1996a). Because seabirds whose populations declined had diets consisting primarily of age-1 pollock (Springer 1992, Decker et al. 1995, Hunt et al. 1996b, 2002a), and the proportion of these declined in their diets, it was hypothesized that declines in the abundance of age-1 pollock and other forage fishes reduced the foraging success and, consequently, reproductive

success and population size of kittiwakes and murrelets at the Pribilof Islands (Hunt and Byrd 1999). A shift from use of age-1 to age-0 pollock during this period (Hunt et al., 1996b) would have been particularly challenging, as age-0 pollock are both smaller (about 2 g) than age-1 pollock in spring (about 19 g) and of lower energy density (about 2% fat vs. about 5% fat) (Iverson et al., 2002).

The decreases in at least black-legged kittiwakes may be related to an increase in adult pollock on the shelf and competition with these fish for age-1 pollock and other forage fish. The biomass of adult (age 3+) walleye pollock on the Bering Sea shelf increased during 1976–1999 ( $r^2 = 0.30$ ,  $P = 0.006$ ,  $n = 24$ ), with the steepest increase in their biomass occurred during 1979–1982 (Vlietstra et al., 2003). The annual variability in reproductive success in black-legged kittiwakes was negatively correlated with the biomass of adult walleye pollock on the Bering Sea shelf in the same year ( $r = -0.68$ ,  $p < 0.001$ ) (Livingston et al., 1999; Hunt and Stabeno, 2002; Vlietstra et al., 2003). There was no correlation between reproductive success in red-legged kittiwakes ( $r = -0.35$ ,  $p = 0.160$ ), common murrelets ( $r = 0.18$ ,  $p = 0.626$ ), or thick-billed murrelets (St. Paul:  $r = -0.419$ ,  $p = 0.136$ ; St. George:  $r = 0.19$ ,  $p = 0.440$ ) and the biomass of adult walleye pollock on the shelf (Vlietstra et al., 2003)

### **Prey Availability and Seabird Mortality at the Pribilof Islands**

Although Vlietstra et al. (2003) found little correlation between juvenile pollock abundance on bottom trawl surveys and seabird reproductive success on an interannual basis, changes in the availability young pollock (and possibly other prey) to seabirds during the late 1970s probably *did* result in poor foraging conditions for seabirds. Inter-colony comparisons suggest that seabird productivity at the Pribilof Islands is limited by the availability of food. For example, since the mid-1970s, the long-term (1976–1999) mean reproductive success of black-legged kittiwakes on St. Paul Island is 0.28 chicks/nest/year, and on St. George Island, it is 0.22 chicks/nest/year. In contrast, mean reproductive success over the same period for black-legged kittiwakes at Cape Lisburne, northern Bering Sea, is 0.80 chicks/nest/year, and in some areas, black-

legged kittiwakes raise two or more chicks per year. At the Pribilofs, kittiwakes raise at most one (Dragoo et al. 2001). Likewise, reproductive success is also low among murrelets at the Pribilof Islands relative to murrelets at other colonies in the North Pacific. Since the mid 1970s, common murrelets on St. Paul Island fledged an average of 0.53 chicks/nest site/year, and, on St. George Island, the average is 0.52 chicks/nest site/year. In contrast, common murrelets at Cape Thompson, in the northern Bering Sea, fledged an average of 0.70 chicks/nest site/year. At Kodiak Island (E. Amatuli), in the Gulf of Alaska, they fledged an average of 0.71 chicks/nest site/year (Dragoo et al. 2001). The ability of kittiwakes and murrelets at other colonies to consistently raise more young than they do at the Pribilof Islands suggests that foraging conditions around the Pribilofs are poor relative to other regions in the Bering Sea.

If prey availability to seabirds at the Pribilof Islands is relatively low during the breeding period, as suggested in the preceding discussion, reproduction may be an especially stressful time for Pribilof seabirds. Kitaysky and colleagues (1999a,b; 2000, 2001) demonstrated the exacerbating effect that a limited food supply can have on acute stress to seabirds during the reproductive phase. Kitaysky et al. (1999b) found that adult black-legged kittiwakes breeding at a colony in Cook Inlet, Alaska, had elevated corticosterone levels during reproduction, and that stress levels were negatively associated with food availability near the breeding colony (Kitaysky et al. 1999b). In addition, corticosterone concentrations in birds increased throughout the breeding season, indicating that stress can become increasingly severe for seabirds as they progress from egg-laying to the chick-rearing period (Kitaysky et al., 1999b). Using captive kittiwakes, Kitaysky et al. (1999a, 2001a) showed that food scarcity and food of poor nutritional quality increased circulating concentrations of a stress-response hormone, corticosterone, in black-legged and red-legged kittiwake chicks. Corticosterone indicates catabolic activity of metabolic pathways, signaling a reliance by seabirds on endogenous energy reserves.

Increasing levels of corticosterone in plasma may affect not only the behaviors of the birds involved, but also their probability of survival. Kitaysky et al., (2000) showed experimentally that adult black-legged kittiwakes with elevated levels of corticosterone had significantly lower probability of returning to their colonies over the two years subsequent to implantation of corticosterone than did birds with sham operations. These results support the hypothesis that when adults have to work hard to raise chicks, either because food may be in short supply (Kitaysky et al., 2000) or because the begging of hungry chicks forces them to work harder (Kitaysky et al., 2001b), adult survival may be compromised (Golet et al., 1998; Golet and Irons, 1999; Kitaysky et al., 2000b). In the laboratory, Kitaysky et al., (2003) showed that young red-legged kittiwake chicks with elevated levels of corticosterone resulting from implants were less able at learning to find food than were birds that were sham-operated. Here the implication is that young kittiwakes that have had less food than optimal while growing may have difficulty learning to feed themselves once fledged and independent. These birds would be exceptionally vulnerable to starvation during the first few months of independence. There is considerable evidence that young seabirds have a more difficult time catching prey than adults. Learning deficits imposed by lack of adequate feeding in the nest and the consequent high stress levels would exacerbate this situation.

Thus, mortality among seabirds at the Pribilof Islands may have varied with the abundance of prey near the islands and may have been most severe during the post-breeding period, when deaths were difficult to observe. By that time, adults would have spent a considerable amount of energy on reproduction and fledglings must learn to forage for themselves or perish. Storm activity is also greater during the post-breeding period (i.e., autumn), and inclement weather can have both direct and indirect implications for chick survivorship and adult mortality, as stormy weather impairs foraging by some seabirds. At the Pribilof Islands, Braun and Hunt (1983) found that, in black-legged kittiwakes on St. Paul Island, chick loss due to ejection of second-hatched young by an older sibling happened more often than expected by chance during

inclement weather. Work by others has shown that these behaviors result from the older chick being unable to satisfy its nutritional needs (Mock et al. 1990, Irons, 1992).

Inability to obtain adequate prey has also affected migrant, non-breeding species in the Bering Sea. Each year millions of short-tailed shearwaters (*Puffinus tenuirostris*) migrate from Australia to the Bering Sea to forage over the inner shelf. The evolution of this annual trans-equatorial migration implies that usually extraordinary amounts of prey must be readily available to these birds in the Bering Sea. During summer and fall of 1997, possibly as many as 10% of the 16 million short-tailed shearwaters present in the southeastern Bering Sea died (Vance et al. 1998; Baduini et al, 2001a, 2001b). Between 1 August and 1 September, reports of moribund and dead shearwaters came from both sides of the Alaska Peninsula, the Aleutian Islands as far west as Adak, Bristol Bay, the Bering Sea north to Cape Anadyr (Russia), and the eastern Chukchi Sea. At least several hundred thousand shearwaters washed ashore. Compared with previous large-scale seabird die offs in the Bering Sea (Bailey & Davenport 1972; Oka & Maruyama 1986; Hatch 1987; Piatt & Van Pelt 1997), the 1997 event in the southeastern Bering Sea was probably one or two orders of magnitude larger ( $10^5$  -  $10^6$  vs.  $10^4$  birds dying). Apparently they were unable to obtain sufficient of the usual prey, euphausiids, in 1997. In 1998, shearwaters were again emaciated, but few died, perhaps because the summer was windier, and they needed less energy to fly between places where food was obtainable. Their diets also shifted, with a greater proportion of fish and zooplankton being taken than was true in the same region in earlier years (Hunt et al., 2002c).

In spring 1998, shearwaters had lower body masses than in spring 1997. Because the seasonal progression of stratification, bloom, and euphausiid mating occurred earlier in 1997, foraging for shearwaters arriving in May 1997 was apparently good, and birds were fat in June. In contrast, in 1998 the seasonal progression was late, and euphausiids were only beginning to form mating swarms in June. Consequently, shearwaters did not put on the level of fat reserves seen in June 1997.

Perhaps because euphausiid mating came early in 1997, by late July few near-surface swarms of mating euphausiids were available to shearwaters. If we are correct in our assessment that euphausiid maturation was delayed in 1999 and that the biomass of euphausiids was large compared to 1997, then swarms should have been available late into the summer and late-season body mass of shearwaters should be high. In the cold spring of 1999, euphausiids were abundant well into summer (Coyle and Pinchuk, 2002a), and shearwaters gained mass through the summer. Thus, it appears that temperature, through its effects on the timing of the availability of prey, may have extremely important impacts on the survival of top predators.

### **What controls recruitment of walleye pollock?**

Walleye pollock dominate the biomass of the eastern Bering Sea ecosystem and the trophic guilds in which they participate (Schumacher et al., In press). They are harvested in greater quantity than any other species of fish or shellfish, and made up > 65% of the total groundfish biomass harvested during the 1980s (Napp et al., 2000). Thus, in any examination of the ecosystems of the eastern Bering Sea, factors that influence the abundance, distribution and ecology of walleye pollock are of key importance for understanding the function of the ecosystem as a whole.

Walleye pollock were not always so abundant. In the early 1960s their populations were small by comparison with those of the 1980s and 1990s. The reason for their initial increase in the mid- to late-1960s is not known, but one hypothesis is that the increase was a result of a trophic cascade that occurred when large, plankton-eating whales and pacific ocean perch (*Sabastes alutus*) were harvested to “commercial extinction” in the late 1960s and early 1970s thereby releasing vast amounts of zooplankton from consumption (Merrick, 1997; NRC, 1996). This newly available zooplankton resource then fuelled the expansion of the pollock population. The removal of the whales may also have released age-0 and age-1 pollock from predation pressure, as fin (*Balaenoptera physalus*) and minke (*B. acutorostrata*) whales would have most likely consumed young pollock along with zooplankton.

During the 1970s and early 1980s, the biomass of walleye pollock first declined and then increased explosively. In the mid- 1970s excessively high harvest rates imposed by the foreign fishing fleets caused the stock to decline. This foreign fishery ceased as of the establishment of the 200-mile Exclusive Economic Zone in 1977. In 1978, pollock produced an enormous year-class that numbered 63.5 billion age-1 fish in 1979 (NMFS, 2002). As these fish grew to adult size, their biomass came to dominate the eastern Bering Sea ecosystem. The reasons for this second spurt in population growth are not known. It is possible that this was a continuation of the ecological release of the pollock population subsequent to the removal of the whales and fish with which they are believed to have competed for prey. An alternative, but not mutually exclusive hypothesis for this population growth is that the strong year-classes of 1977 and 1978 were the result of the 1976/77 regime shift that resulted in earlier ice retreat and warmer water temperatures in spring, conditions favorable for zooplankton production that would have provided abundant prey resources for growing pollock (Francis et al., 1998; Hunt et al., 2002a). Indeed, there is a general acknowledgment that warm years are conducive to enhanced pollock recruitment (Bulatov, 1995; Hollowed and Wooster, 1995; Bailey et al., 1995, 1996; Quinn and Niebauer, 1995; Walsh and McRoy, 1986; Blood, 2002). The cessation of the foreign fishery did not cause the exceptional year class of 1978. The increase was driven by exceptional early survival and recruitment. Had the fishery continued into the mid-1980s, there might not have been as great a growth in pollock biomass, but the 1978 year-class still would have been exceptional.

Throughout the 1980s, the biomass of walleye pollock in the eastern Bering Sea remained high, though between 1990 and 1998 there was a substantial decline (Figure 32) (NMFS , 2002). During the 1980s, there were 3 strong year-classes, and during the 1990s there were 2 (Table 6; Figure 32). It is these strong year classes, as they matured, that sustained the biomass of the stock and its ability of support fisheries.

Understanding of the determination of year-class strength is a key to understanding much of what occurs in the southeastern Bering Sea marine ecosystem.

Several hypotheses that have been put forward as explanations for variation in pollock year-class strength (Table 7). They can be divided into four groups: Group A focuses on the importance of primary or secondary production for the survival of young pollock; Group B focuses on the importance of cannibalism and mechanisms that result in the separation of adult and juvenile pollock; Group C is an hypothesis that combines variation in risk of predation with variation in the abundance of zooplankton, and Group D is an hypothesis that relates survival of young pollock to variation in the amount of turbulence when feeding. Ladd et al. (ms) have modelled the amount of entrainment expected in each summer over the past decades and a preliminary comparison of their results with the distribution of strong and weak year-classes in Table 6 suggests that variations in primary production forced by summer entrainment of nutrient is not likely a good predictor of year-class strength in pollock. The abundance of small shelf copepods, does vary significantly with integrated water temperature (Tables 3 and 4), and if prey is a limiting factor in pollock year-class strength, then this remains a viable hypothesis. Cannibalism is recognized as a major source of mortality for young pollock (Dwyer et al., 1987; Wespestad, 1994; Balykin, 1996; Livingston and Lang, 1996; Wespestad and Quinn, 1996; Livingston and Methot, 1998; Lang et al., 2000; Livingston and Juardo-Molina, 2000), and mechanisms for separating vulnerable individuals from cannibalistic older fish, such as on-shelf advection (Wespestad et al., 2000), are likely to be of importance regardless of what other factors are at play. For instance, Cooney et al. (2001) found that in Prince William Sound adult pollock would feed on large *Neocalanus* copepods in spring if the copepods were abundant, but if not, the adult pollock would prey upon juvenile salmon and herring. Finally, variations in the amount of turbulence may influence foraging success of larval pollock be either enhancing the encounter rate with food particles at intermediate rates of turbulence, or depressing foraging success with either too much turbulence or insufficient turbulence (Megrey and Hinckley, 2001). Most of these hypotheses remain to be tested in the field, and it is

likely that some combination of them together will provide the key to understanding the occurrence of strong year-classes.

It is also likely that, as pollock populations wax and wane, the mechanisms responsible for population control will shift. Hunt et al. (2002a) hypothesized that under environmental conditions in which zooplankton were scarce, such as during a prolonged series of cold springs, walleye pollock populations would be controlled from the bottom-up because there would be insufficient zooplankton for larval or juvenile pollock (Figure 33). In contrast, in years when pollock populations were large, cannibalism would exert strong top-down control on the population, as it apparently does at present. In periods under top-down control, large year-classes could still occur, but to do so there would have to be some mechanism to promote separation of the young fish from the adults (e.g., Hypotheses 3 and 4, Table 7). When there is a switch from a period of years with cold springs to warm springs, one can expect large year classes, as there should be plentiful zooplankton and low risk of cannibalism, while during a switch from a prolonged period of warm springs to cold springs, year-class strength might be expected to be low because of both bottom-up and top-down effects.

The distributions and abundances of adult and juvenile pollock (age-0 and age-1) have important impacts on other elements of the eastern Bering Sea ecosystem. Juvenile pollock are a major source of prey for not only large piscivorous fishes such as adult pollock, Pacific cod and arrowtooth flounder, they are also staples in the diets of several species of marine birds and pinnipeds, such as northern fur seal, harbor seal, and Steller sea lions (see above). There is evidence for a negative relationship between the reproductive success of black-legged kittiwakes on St George Island and the biomass of adult pollock on the eastern Bering Sea shelf (Livingston et al., 1999; Hunt and Stabeno, 2002; Vlietstra et al., 2003). During the period of population decline in the 1990s, northern fur seals at St. Paul island in the Pribilof islands decreased their use of age-0 and age-1 pollock, sand lance, and slope squids, but increased their use of

age-3, -4, -5 pollock. It is difficult to know the causes of these shifts in fur seal diets, but it appears that the shift in diet is not beneficial to these pinnipeds.

The distribution of age-1 pollock may be affected by not only water temperature (Hypothesis 3, Table 7), but also their exposure to predation, either through removals, or through behavioral responses to the presence of predators (Sogard and Olla, 1993). Displacements of juvenile pollock from surface waters will make them unavailable to predators such as kittiwakes that are restricted in their foraging to the very surface of the water column, while displacements in the horizontal from the vicinity of rookeries and colonies will diminish their value to central place foragers as a function of the distance travelled from the central place. After the biomass of adult pollock increased in the vicinity of the Pribilof Islands, the biomass of age-1 pollock there dropped precipitously (Hunt et al., 2002a, Fig12). There remains a need to understand the role that adult pollock play in determining not only the abundance, but also the distribution of forage fishes, including age-1 pollock, and their availability to upper trophic level predators including marine mammals and seabirds.

### **Summary of Advances**

Many of the important advances in our understanding of the coupling of regional atmospheric processes, physical processes in the Bering Sea, and biological responses to these forcing mechanisms have been the subjects of recent review (Stabeno et al., 2001; Overland et al., 2002; Schumacher et al., In Press). Among these advances in our knowledge are:

Marine Climate:

- 1) A new understanding of the importance of warm-season climate (Overland et al., 2001; Bond and Adams, 2002);

Basin and Slope Waters:

- 1) An improved understanding of the function of the Aleutian North Slope Current and the Bering Slope Current (Reed and Stabeno, 1999; Stabeno et al., 1999; Johnson et al., submitted), and their potential importance to fish (Reed, 1995);
- 2) Documentation of eddies over the slope (Schumacher and Stabeno, 1994; Okkonen, 2001) and along the outer shelf (Reed, 1998) and their importance as habitats for fish and for the transport of small pollock onto the shelf (Schumacher and Stabeno, 1994).
- 3) Development of a model that resolves the eddies and suggests that the source of the eddies present in the southeastern Bering Sea basin originate in the Bering slope current (Hermann et al. 2002).

#### Shelf Waters:

- 1) Discovery of mean flows across the shelf (Reed and Stabeno, 1996; Reed, 1998), which for short intervals in summer can result in replenishment of salts and transport of oceanic zooplankton to the Inner Front (Stabeno et al., 2001; Coyle and Pinchuk, 2002b);
- 2) A clarification of the role of the timing of ice retreat for the timing and fate of the spring bloom (Stabeno et al., 2001; Hunt et al., 2002a; Hunt and Stabeno, 2002);
- 3) Understanding of the relative importance of winter and warm season conditions for determining the timing and strength of spring stratification and the potential for summer mixing across the pycnocline (Ladd et al., ms);
- 4) An understanding of the temporal and spatial variability of the Inner Front and its role in nutrient fluxes from depth (Kachel et al., 2002);
- 5) A greater appreciation for the role of water temperature in the distribution of fish, especially pollock (Ohtani and Azumaya, 1995; Wyllie-Echeverria, 1996; Wyllie-Echeverria and Wooster, 1998), the hatching time of pollock eggs (Blood, 2002), and the population dynamics of zooplankton (Coyle and Pinchuk, 2002a, b);

- 6) A greater appreciation for the role of episodic events in structuring the ecosystem;
- 7) A new hypothesis concerning the role of wind-forced transport for the survival of juvenile walleye pollock (Wespestad et al., 2000).

#### Biological Components of the Ecosystem

- 1) Evidence that the net annual primary production in the northern Bering Sea may be declining (Schell, 2000; Grebmeier and Cooper, 1994; Grebmeier and Dunton, 2000), whereas in the southeastern Bering Sea, increases in standing stocks of fish and non-crab benthic invertebrates do not suggest a decline in production (Connors et al, 2002).
- 2) Commencement of research on the role of nanno- and micro-plankton, including coccolithophores, in the marine ecosystem of the southeastern Bering Sea (Howell-Kubler et al. 1996; Olson and Strom, 2002).
- 3) Re-examination of the Oshoro Maru zooplankton time-series shows no evidence for a significant decline in zooplankton biomass within samples stratified by the domain in which they were collected (Hunt et al., 2002; Napp et al., 2002).
- 4) Immense increases in the biomass of large jellyfish and then a rapid decline to levels seen in the early 1980s (Brodeur et al., 1999, 2002).
- 5) Observations of responses of crustacean zooplankton to variations in water temperature, including apparent shifts in the timing of breeding in shelf euphausiids (Coyle and Pinchuk, 2002a), and the abundance of small shelf copepods (Coyle and Pinchuk, 2002a; Napp et al, 2002)
- 6) Determination that large cetaceans are now encountered more frequently in shelf waters than in the 1970s, and that substantial numbers are present over the shelf (Baretta and Hunt 1994; Moore, 2000).
- 7) Progress in defining multiple hypotheses concerning the control of year-class strength in walleye pollock (Indices Working Group Report, this volume).

- 8) Improved understanding of the foraging habits and diets of marine birds and northern fur seals, and Steller sea lions at the Pribilof Islands, and the potential relationships between the abundance of adult pollock over the shelf and the reproductive success of the black-legged kittiwake at the Pribilof Islands (Vlietstra et al., 2003; Sinclair et al., In prep).
- 9) Evidence the sub-lethal food stress during the breeding season can lead to elevated levels of corticosteroids, and possibly elevated levels of post-breeding season mortality in seabirds (Kitaysky et al., 199b, 2001a, 2002)
- 10) Development of a mechanistic explanation of how variations in climate could cause shifts in the mechanisms controlling walleye pollock populations (Oscillating Control Hypothesis, Hunt et al., 2002a).

### **Areas of Research Requiring More Attention**

Although a great deal of progress in understanding the forcing mechanisms and the ecosystem responses of the southeastern Bering Sea has been made since the early 1990s, there are still many unanswered questions that demand attention if we are to provide good stewardship for the resources of the Bering Sea shelf. Answers to these questions go beyond academic value; we need to understand the processes and mechanisms that determine how changes in global climate and the associated physical forcing affect the flow of energy to upper trophic level organisms including commercially valuable fish and shellfish, marine mammals and seabirds. Below listed a number of questions that require attention:

- 1) How does atmospheric forcing affect cross-shelf flow? Do the mechanisms whereby climate affects cross-shelf flow vary seasonally?
- 2) What is the relative importance of nutrients provided by wintertime on-shelf advection and nutrients that have been remineralized by benthic processes? Are the estimates provided by Sambrotto et al. (1986) still useful?

- 3) How does the pattern of ice-melt back affect the timing and type of phytoplankton bloom. Do these mechanisms work in similar ways in the southern and northern Bering Sea?
- 4) Do open-water blooms occur in the northern Bering Sea, and if so, are they as important as the ice-related blooms?
- 5) Are ice-related blooms shorter and more intense than open water blooms? When both are potentially present, which is likely to fix more carbon?
- 6) Has primary production decreased over the southeastern Bering Sea shelf?
- 7) For the cold-adapted species of the Bering Sea, are zooplankton growth and production more sensitive to temperature than phytoplankton productivity, as is assumed in the hypothesis of Walsh and McRoy (1986)?
- 8) Does significantly more phytoplankton sink to the bottom in cold years than in warm years, as hypothesized by Walsh and McRoy (1986)?
- 9) How do the nanno- and micro-plankton affect the flow of energy in the southeastern Bering Sea? Are they directly or indirectly important prey for either larval or juvenile walleye pollock?
- 10) How do coccolithophore affect energy flow in the southeastern bering Sea, and why did they suddenly bloom, and as suddenly disappear after blooming consecutively for five years?
- 11) How do plankton predators such as chaetognaths, ctenophores, and larger jellyfish affect the flow of energy in the eastern Bering Sea? Why did large jellyfish increase and then crash? Are they significant competitors or predators on juvenile pollock and other forage fish?
- 12) How does temperature affect the role that chaetognaths, ctenophores and larger jellyfish as predators of crustacean zooplankton?
- 13) How can one account for the simultaneous increase of round-fish, flatfish and non-crab invertebrates over the shelf of the southeastern Bering Sea (Connors et al., 2002)?

- 14) How variable is the abundance of mesopelagic fish and squids in the eastern Bering Sea, and what are the ramifications of this variability for their use by commercially valuable fish, marine mammals and seabirds?
- 15) What factor or factors control year-class size in walleye pollock? There is a need for systematic testing of the most promising hypotheses, such as those listed in Table 7.
- 16) There is a need to test the underlying assumptions and predictions of the Oscillating Control Hypothesis (Hunt et al., 2002a). In particular, there is a need to determine if zooplankton grazers are closely coupled with phytoplankton in any season, and if this coupling affects their availability to larval or juvenile pollock. Likewise, it is important to determine whether larval or juvenile pollock are closely coupled to zooplankton in any season.
- 17) Determination as to whether chronic food stress in pinnipeds leads to elevated corticosteroid levels and decreased post-breeding survival.

## **Acknowledgements**

Many scientists provided text and figures that have been incorporated into this report. Included are Nick Bond, Ric Brodeur, Liz Conners, Susan Henrichs, Anne Hollowed, Jim Overland, Beth Sinclair, Phyllis Stabeno, and Al Tyler. Financial support for this review was provided in part by the NOAA Coastal Ocean Program, Southeast Bering Sea Carrying Capacity program.

## References Cited-- Plus lots of extras that need to be eliminated!

Alexander, V., Cooney, T., 1979. A quantitative study of the phytoplankton from the eastern Bering Sea. In: Environmental Assessment of the Alaska Continental Shelf. NOAA/OCSEAP, Annual Report.

Anderson, P.J., Piatt, J.F., 1999. Community reorganization in the Gulf of Alaska following ocean climate regime shift. Marine Ecology Progress Series 189, 117-123.

Anonymous, 1993. Is It Food? Addressing marine mammal and seabird declines. Workshop Summary. Alaska Sea Grant Report 93-01.

Antonelis, G.A., Sinclair, E.H., Ream, R.R., Robson, B.W., 1997. Inter-island variation in the diet of female northern fur seals (*Callorhinus ursinus*) in the Bering Sea. Journal Zoological Society London 242, 435-451.

Aydin, K.Y., Lapko, V.V., Radchenko, V.I., Livingston, P.A., In Press. A comparison of the eastern and western Bering Sea shelf/slope ecosystems through the use of mass-balance food web models. NOAA Technical Memo ###. 76p.

Baduini, C.L., Hyrenbach, K.D., Coyle, K.O., Pinchuk, A., Mendenhall, V., and Hunt, Jr., G.L., Jr., 2001. Mass mortality of short-tailed shearwaters in the eastern Bering Sea during summer 1997. *Fisheries Oceanography* 10, 117-130.

Baduini, C.L., Lovvorn, J.R., Hunt, G.L., Jr. 2002. Determining the nutritional condition of short-tailed shearwaters (*Puffinus tenuirostris*): implications for migratory flight ranges and starvation events. Marine Ecology Progress Series 222: 265-277.

Bailey, K.M., 2000. Shifting control of recruitment of walleye pollock *Theragra chalcogramma* after a major climatic and ecosystem change. *Marine Ecology Progress Series* 198, 215-224.

Bailey, K.M., Brown, A.L., Yoklavich, M.M., Mier, K.L., 1996. Interannual variability of growth of larval and juvenile walleye pollock in the western Gulf of Alaska. *Fisheries Oceanography* 5 (Supplement 1), 137-147.

Bailey, K.M., Canino, M.F., Napp, J.M., Spring, S.M., Brown, A.L., 1995. Contrasting years of prey levels, feeding condition and mortality of larval walleye pollock (*Theragra chalcogramma*) in the western Gulf of Alaska. *Marine Ecology Progress Series* 119, 11-13.

Bailey, K.M., Powers, D.M., Quattro, J.M., Villa, G., Nishimura, A., Traynor, J.J., Walters, G., 1999. Population ecology and structural dynamics of walleye pollock (*Theragra chalcogramma*). In: Loughlin, T.R., Ohtani, K. (Eds.), *Dynamics of the Bering Sea: A Summary of Physical, Chemical, and Biological Characteristics, and a Synopsis of Research on the Bering Sea*. North Pacific Marine Science Organization (PICES), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, USA, pp. 581-614.

Baird, P.A., 1990. Influence of abiotic factors and prey distribution on diet and reproductive success of three seabird species in Alaska. *Ornis Scandinavica* 21, 224-235.

Bakkala, R.G., 1993. Structure and historical changes in the groundfish complex of the eastern Bering Sea. NOAA Technical Report NMFS 114, 91pp.

Balch, W.M., Holligan, P.M., Ackleson, S.G. and Voss, K.J. (1991) Biological and optical properties of mesoscale coccolithophore blooms in the Gulf of Maine. *Limnol. Oceanogr.* **36**:629-643.

Balch, W.M., Holligan, P.M., Kilpatrick, K.A. 1992. Calcification, photosynthesis and growth of the bloom-forming coccolithophore, *Emiliana huxleyi*. Cont. Shelf. Res. 12: 1353-1374.

Balykin, P.A., 1996. Dynamics and abundance of western Bering Sea walleye pollock. In: Mathisen, O.A., Coyle, K.O. (Eds.), Ecology of the Bering Sea: A review of Russian literature. Univ. Alaska Sea Grant Report 96-01, Fairbanks, Alaska, pp. 177-182

Baretta, L., Hunt, G.L., Jr., 1994. Changes in the numbers of cetaceans near the Pribilof Islands, Bering Sea, between 1975-78 and 1987-89. Arctic 47, 321-326.

Barnston, A.G., Livezey, R.E. 1987. Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. Monthly Weather Review 115: 1083-1126.

Beamish, R.J., 1993. Climate and exceptional fish production off the west coast of North America. Canadian Journal of Fisheries and Aquatic Science 50, 2270-2291.

Beamish, R.J., Leask, K.D., Ivanov, O.A., Balanov, A.A., Orlov, A.M., Sinclair, B. 1999. The ecology, distribution, and abundance of midwater fishes of the Subarctic Pacific gyres. Progress in Oceanography 43: 399-442.

Beamish, R.J., Noakes, D.J., McFarlane, G.A., Klyashtorin, L., Ivanov, V.V., Kurashov, V., 1999. The regime concept and natural trends in the production of Pacific salmon. Canadian Journal of Fisheries and Aquatic Science 56, 516-526.

Blood, D.M. 2002. Low-temperature incubation of walleye pollock (*Theragra chalcogramma*) eggs from the southeastern Bering Sea shelf and Shelikof Strait, Gulf of Alaska. Deep-Sea Research Part II 49: 6095-6108.

Bond, N.A., Adams, J.M., (2002) Atmospheric forcing of the southeast Bering Sea shelf during 1995-1999 in the context of a 40-year historical record. *Deep-Sea Research Part II* 49: 5869-5887.

Braun, B.M. and G.L. Hunt. 1983. Brood reduction in Black-legged Kittiwakes. *Auk* 100: 469-476.

Brodeur, R.D., Mills, C.E., Overland, J.E., Walters, G.E., Schumacher, J.D., 1999a Evidence for a substantial increase in gelatinous zooplankton in the Bering Sea, with possible links to climate change. *Fisheries Oceanography* 8, 296-306.

Brodeur, R.D., Sugisaki, H., & Hunt, G.L. Jr. 2002. Increases in jellyfish biomass in the Bering Sea: implications for the ecosystem. *Marine Ecology Progress Series* 233: 89-103.

Brodeur, R.D., Wilson, M.T., Walters, G.E., Melnikov, I.V., 1999b Forage fish in the Bering Sea: distribution, species associations and biomass trends In: Loughlin, T.R., Ohtani, K. (Eds.), *Dynamics of the Bering Sea: A Summary of Physical, Chemical, and Biological Characteristics, and a Synopsis of Research on the Bering Sea*. University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, USA, pp. 509-536.

Brodeur, R.D., Wilson, M.T., Ciannelli, L., 2000. Spatial and temporal variability in feeding and condition of age-0 walleye pollock (*Theragra chalcogramma*) in frontal regions of the Bering Sea. *ICES Journal of Marine Science*. 57, 256-264.

Brown, C.W. and Podesta, G.P. (1997) Remote sensing of coccolithophore blooms in the western South Atlantic Ocean. *Remote Sens. Environ.* 60:83-91.

Brown, C.W. and Yoder, J.A. (1993) Blooms of *Emiliania huxleyi* (Prymnesiophyceae) in surface waters of the Nova Scotian shelf and the Grand Bank. *J. Plankton Res.* **15**:1429-1438.

Brueggerman, J.J., Oliver, G.W., Fowler, C., Frost, K., Fay, K., Cowles, C., Schneider, K., Calkins, D., 1987. Marine Mammals. In: Hameedi, M.J. (Ed.), The St. George Basin environment and possible consequences of planned offshore oil and gas development. U.S. Depart. Commerce, U.S. Dept. Interior, Outer Continental Shelf Environmental Assessment Program, Juneau, Alaska, USA, pp. 55-81.

Bulatov, O.A., 1995. Biomass variations of walleye pollock in the Bering Sea in relation to oceanological conditions. In: Beamish, R.J. (Ed.), Climate change and northern fish populations. Canadian Special Publication in Fisheries and Aquatic Science 121, 631-640.

Capriulo, G.M., Sherr, E.B., Sherr, B.F. 1991. Trophic behavior and related community feeding activities of heterotrophic marine protists. In: Reid, P.C., Turley, C.M., Burkil, P.H. (Eds.), Protozoa and their role in marine processes. Springer, Berlin, pp.219-265.

Cavalieri, D.J. and Martin, S. 1994. The contribution of Alaskan, Siberian and Canadian coastal polynyas to the cold halocline layer of the Arctic Ocean. *J. Geophysical Research* 99: 18343-34862 ???

Ciannelli, L., Brodeur, R., Buckley, T.W., 1998. Development and application of a bioenergetics model for juvenile walleye pollock. *Journal of Fish Biology* 52, 879-898.

Coachman, L.K., 1986. Circulation, water masses and fluxes on the southeastern Bering Sea shelf. *Continental Shelf Research* 5, 23-108.

Coachman, L.W., Kinder, T.H., Schumacher, J.D., and Tripp, R.B. 1980. Frontal systems of the southeastern Bering Sea shelf. In: Carstens, T., McClimans, T. (Eds) Stratified flows, second IAHR Symposium, Trondheim, June 1980. Tapir, Trondheim, pp. 917-933.

Conners, M.E., Hollowed, A.B., Brown, E. 2002. Retrospective analysis of Bering Sea bottom trawl surveys: regime shift and ecosystem reorganization. *Progress in Oceanography* 55: 209-222.

Cooney, R.T., Coyle, K.O., 1982. Trophic implications of cross-shelf copepod distributions in the southeastern Bering Sea. *Marine Biology* 70, 187-196.

Cooney, R.T., Allen, J.R., Bishop, M.A., Eslingler, D.L., Kline, T., Norcross, B.L., McRoy, C.P., Milton, J., Olsen, J., Patrick, V., Paul, A.J., Salmon, D., Schell, D., Thomas, G.L., Vaughan, S.L., Willette, T.M. 2001. Ecosystem controls of juvenile pink salmon (*Onchorynchus gorbusha*) and Pacific herring (*Clupea pallasii*) populations in Prince William Sound, Alaska. *Fisheries Oceanography* 10 (Suppl 1): 1-13.

Corkett, C.J., McLaren, I.A., 1978. The biology of *Pseudocalanus*. *Advances in Marine Biology* 15, 1-231.

Coyle, K.O., Chavtur, V.G., Pinchuk, A.I., 1996. Zooplankton of the Bering Sea: a review of the Russian literature. In: Mathisen, O.A., Coyle, K.O. (Eds.), *Ecology of the Bering Sea: A review of Russian literature*. Univ. Alaska Sea Grant Report 96-01, Fairbanks, Alaska, USA, pp. 97-133.

Coyle, K.O., Pinchuk, A.I., 2002a. The abundance and distribution of euphausiids and zero-age pollock on the inner shelf of the southeast Bering Sea relative to the inner front. *Deep-Sea Research Part II* 49: 6009-6030.

Coyle, K.O., Pinchuk, A.I. 2002b. Climate-related differences in zooplankton density and growth on the inner shelf of the southeastern Bering Sea. *Progress in Oceanography* 55: 177-194.

Cullen, J.T., Rosenthal, Y., Falkowski, P.G. 2001. The effect of anthropogenic CO<sub>2</sub> on the carbon isotope composition of marine phytoplankton. *Limnology and Oceanography* 46: 996-998.

Dagg, M.J., Clarke, M.E., Nishiyama, T., Smith, S.L., 1984. Production and standing stock of copepod nauplii, food items for larvae of the walleye pollock *Theragra chalcogramma* in the southeastern Bering Sea. *Marine Ecology Progress Series* 19, 7-16.

Dagg, M.J., Vidal, J., Whitley, T.E., Iverson, R.L., Goering, J.J., 1982. The feeding, respiration, and excretion of zooplankton in the Bering Sea during a spring bloom. *Deep-Sea Research* 29, 45-63.

Decker, M.B., Hunt, G.L., Jr., 1996. Foraging by murre ( *Uria* spp.) at the tidal front surrounding the Pribilof Islands, Alaska. *Marine Ecology Progress Series* 139, 1-10.

Decker, M.B., Hunt, G.L., Jr., Byrd, G.V., Jr., 1995. The relationships among sea surface temperature, the abundance of juvenile walleye pollock (*Theragra chalcogramma*), and the reproductive performance and diets of seabirds at the Pribilof Islands, southeastern Bering Sea. In: Beamish, R.J. (Ed.) *Climate change and northern fish populations*. Canadian Special Publication in Fisheries and Aquatic Science 121: 425-437.

Dragoo, D.E., Byrd, G.V., Irons, D.B., 2000. Breeding status and population trends of seabirds in Alaska in 1999. U.S. Fish and Wildlife Service Report AMNWR 2000/02.

Dragoo, D. E., Byrd, G.V., and Irons, D.B. 2001. Breeding status, population trends and diets of seabirds in Alaska, 2000. U.S. Fish and Wildlife Service Report AMNWR 01/07.

Dugdale, R.C., Goering, J.J., 1967. Uptake of new and regenerated forms of nitrogen in primary productivity. *Limnology and Oceanography* 12, 196-206.

Dussart, B.M. 1965. Les différentes catégories de plancton. *Hydrobiologica* 26: 72-74.

Dwyer, D.A., Bailey, K.M., Livingston, P.A., 1987. Feeding habits and daily ration of walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea, with special reference to cannibalism. *Canadian Journal of Fisheries and Aquatic Science* 44, 1972-1984.

Ebbesmeyer, C.C., Cayan, D.R., Milan, D.R., Nichols, F.H., Peterson, D.H., Redmond, K.T., 1991. 1976 step in the Pacific climate: forty environmental changes between 1968-1975 and 1977-1984. In: Betancourt, J.L., Tharp, V.L. (Eds.), *Proceedings of the Seventh Annual Climate (PACLIM) Workshop*, April 1990. . California Department of Water Resources, Interagency Ecological Studies Program Technical Report 26, pp. 115-126.

Eslinger, D.L., Iverson, R.L., 2001. The effects of convective and wind-driven mixing on spring phytoplankton dynamics in the southeastern Bering Sea middle shelf domain. *Continental Shelf Research* 21, 627-650.

Fernandez et al., 1994

Flint, M.V., Sukhanova, I.N., Kopylov, A.I., Poyarkov, S.G. and Whitley, T.E. 2002. Plankton distribution associated with frontal zones in the vicinity of the Pribilof Islands. *Deep Sea Research Part II*. In Press

Francis, R.C., Hare, S.R., Hollowed, A.B., Wooster, W.S., 1998. Effects of interdecadal climate variability on the oceanic ecosystems of the NE Pacific. *Fisheries Oceanography* 7, 1-21.

Gervais, F., Riebesell, U., 2001. Effect of phosphorus limitation on elemental composition and stable carbon isotope fractionation in a marine diatom growing under different CO<sub>2</sub> concentrations. *Limnology and Oceanography* 46, 497-504.

Golet, G. H., and Irons, D.B. 1999. Raising young reduces body condition and fat stores in black-legged kittiwakes. *Oecologia* 120: 530-538.

Golet, G.H., Irons, D.B. and Estes, J.A. 1998. Survival costs of chick rearing in black-legged kittiwakes. *Journal of Animal Ecology* 67: 827-841.

Gower, J.R.F. 1997. Bright plankton blooms on the west coast of North America observed with AVHRR imagery Pp. 25-41, In: Kahru, M & Brown, C.W. (eds.) *Monitoring algal blooms: New techniques for detecting large-scale environmental change*. Landes Bioscience, Austin, Texas.

Grebmeier, J.M. (1992). Benthic processes on the shallow continental shelf of the northern Bering and Chukchi Seas, pp. 243-251. In: P.A. Nagel (ed) *Results of the Third Joint US-USSR Bering & Chukchi Seas Expedition (BERPAC), Summer 1988*. U.S. Fish and Wildlife Service, Washington, DC.

Grebmeier, J.M., Cooper, L.W. 1994. A decade of benthic research on the continental shelves of the northern Bering and Chukchi Seas: Lessons learned. In: R.H. Meehan, V. Serienko, and G. Weller, Eds. *Bridges of Science between North America and the Russian Far East*. Fairbanks, Alaska: American Association for the Advancement of Science, Arctic Division. P. 87-98.

Grebmeier, J.M., Cooper, L.W. 2002. Benthic processes in the Bering Strait region of the Arctic: Temporal/spatial variability and global change. Eos. Trans. AGU, 83(4), Ocean Sciences Meeting Supplement, Abstract OS32M-01

Grebmeier, J.M., Dunton, K.H. 2000. Benthic processes in the northern Bering/Chukchi Seas: Status and global change. In Huntington, H.P. (Ed.) Impacts of changes in sea ice and other environmental parameters in the Arctic. Report of the Marine Mammal Commission Workshop, 15-17 february 2000, Girdwood Alaska. Available from the Marine Mammal Commission, Bethesda, Maryland.

Hare, S.R., Francis, R.C., 1995. Climate change and salmon production in the northeast Pacific Ocean. In: Beamish, R.J. (Ed.), Climate change and northern fish populations. Canadian Special Publication in Fisheries and Aquatic Science 121, 357-372.

Hare, S. R., Mantua, N.J., 2000. Empirical evidence for North Pacific regime shifts in 1977 and 1989. Progress in Oceanography 47, 103-146.

Hatch, S.A., Byrd, G.V., Jr., Irons, D.B., Hunt, G.L., Jr., 1993. Status and ecology of kittiwakes (*Rissa tridactyla* and *R. brevirostris*) in the North Pacific. In: Vermeer, K, K.T. Briggs, K.H. Morgan, D.C. Siegel-Causey (Eds.), The Status, Ecology and Conservation of Marine Birds of the North Pacific Ocean. Canadian Wildlife Service Special Publication, Ottawa, Canada, pp. 140-153.

Haynes, E.G., Ignell, G., 1983. Effect of temperature on rate of embryonic development of walleye pollock, *Theragra chalcogramma*. Fisheries Bulletin 81, 890-894.

Hillgruber, N., Haldorson, L.J., Paul, A.J., 1995. Feeding selectivity of larval walleye pollock *Theragra chalcogramma* in the oceanic domain of the Bering Sea. Marine Ecology Progress Series 120, 1-10.

Hirons, A.C., Schell, D.M., Finny, B.P., 2001. Temporal records of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  in North Pacific pinnipeds: inferences regarding environmental change and diet. *Oecologia* 129, 591-601.

Holligan, P.M., Harris, R.P., Newell, R.C., Harbour, D.S., Head, R.N., Linley, E.A.S., Lucas, M.I., Tranter, P.R.G., Weekley, C.M., 1984. Vertical distribution and partitioning of organic carbon in mixed frontal and stratified water of the English Channel. Marine Ecology Progress Series 17, 201-213.

Holligan, P.M., Viollier, M., Harbour, D.S., Camus, P. and Champagne-Philippe, M. (1983) Satellite and ship studies of coccolithophore production along a continental shelf edge. *Nature* **304**:339-342.

Hollowed, A.B., Wooster, W.W. 1992. Variability of winter ocean conditions and strong year classes of northeast Pacific groundfish. ICES Mar Sci Symposium 195: 433-444.

Hollowed, A.B., Wooster, W.W. 1995. Decadal-scale variations in the eastern Subarctic Pacific: II Responses of Northeast Pacific fish stocks. Pp. 373-385 in Climate change and northern Fish Populations, Beamish, R.J., ed. Canadian Journal of Fisheries and Aquatic Sciences 121 (Special publication).

Hollowed, A.B., Hare, S.R., Wooster, W.S., 2001. Pacific-Basin climate variability and patterns of northeast Pacific marine fish production. Progress in Oceanography 49, 257-282.

Hood, D.W. 1999. PROBES: Processes and Resources of the eastern Bering Sea shelf. In: Loughlin, T.R., Otahni, K. (Eds.), Dynamics of the Bering Sea. University of Alaska Sea Grant AK-SG-99-03 Fairbanks, pp. 387-407.

Howell-Kubler, A., Lessard, E., Napp, J. 1996. Springtime microprotozoan abundance and biomass in the southeastern Bering Sea and Shelikof Strait, Alaska. *Journal of Plankton Research* 18: 731-745.

Hunt, G.L., Jr., Byrd, G.V., Jr., 1999. Marine bird populations and carrying capacity of the eastern Bering Sea. In: Loughlin, T.R., Ohtani, K. (Eds.), Dynamics of the Bering Sea. Univ. of Alaska Sea Grant AK-SG-99-03, Fairbanks, Alaska, USA, pp.631-650.

Hunt, G.L. and N.M. Harrison. 1990. Foraging habitat and prey taken by Least Auklets at King Island, Alaska. *Mar. Ecol. Prog. Ser.* 65: 141-150.

Hunt, G.L., Jr. and Stabeno, P.J. 2002. Climate change and control of energy flow in the southeastern Bering Sea. *Progress in Oceanography.* 55: 5-22.

Hunt, G.L., Jr., Baduini, C.L., Brodeur, R.D., Coyle, K.O., Kachel, N.B., Napp, J.M., Salo, S.A., Schumacher, J.D., Stabeno, P.J., Stockwell, D.A., Whitledge, T.E., Zeeman, S.I., 1999. The Bering Sea in 1998: A second consecutive year of extreme weather-forced anomalies. *EOS, Transactions of the American Geophysical Union* 80, 561, 565-566.

Hunt, G.L., Jr., Baduini, C., Jahncke, J. 2002b. Diets of short-tailed shearwaters in the southeastern Bering Sea. *Deep Sea Research Part II.* 49: 6147-6156.

Hunt, G.L., Jr., Coyle, K.O., Hoffman, S., Decker, M.B., Flint, E.N., 1996a. Foraging ecology of short-tailed shearwaters near the Pribilof Islands, Bering Sea. *Marine Ecology Progress Series* 141, 1-11.

Hunt, G.L., Jr., Decker, M.B., Kitaysky, A.S., 1996b. Fluctuations in the Bering Sea Ecosystem as reflected in the reproductive ecology and diets of kittiwakes on the Pribilof Islands, 1975 to 1990. In: Greenstreet, S., Tasker, M. (Eds.), *Aquatic Predators and Their Prey*. Blackwell, London, pp. 142-153.

Hunt, G.L., N.M. Harrison and T. Cooney. 1990. Foraging of Least Auklets: The influence of hydrographic structure and prey abundance. *Studies in Avian Biol.* 14: 7-22.

Hunt, G.L., Jr., Kitaysky, A.S., Decker, M.B., Dragoo, D.E., Springer, A.M., 1996c. Changes in the distribution and size of juvenile walleye pollock as indicated by seabird diets at the Pribilof Islands and by bottom trawl surveys in the eastern Bering Sea. In: Brodeur, R.D., Livingston, P.A. Loughlin, T.R., Hollowed, A.B. (Eds.) *Ecology of juvenile walleye pollock, *Theragra chalcogramma**. NOAA Technical Report NMFS 126, 133-147.

Hunt, G.L., Jr., Stabeno, P., Walters, G., Sinclair, E., Brodeur, R.D., Napp, J.M., Bond, N.A. 2002a. Climate change and control of the southeastern Bering Sea pelagic ecosystem. *Deep Sea Research Part II*. 49: 5821-5853.

Huntley, M.E., Lopez, M.D.G., 1992. Temperature-dependent production of marine copepods: a global analysis. *American Naturalist* 140, 201-242.

IARPC (Interagency Arctic Research Planning Committee, 2001. United states Arctic research Plan: Biennial Revision 2002-2006. Office of Polar Programs, National Science Foundation. 84pp.

Iida, T., Saitoh, S.I., Miyamura, T., Toratani, M., Fukushima, H., Shiga, N. 2002. Temporal and spatial variability of coccolithophore blooms in the eastern Bering Sea, 1998-2001. *Progress in Oceanography* 55: 165-175.

Ikeda, T., Motoda, S., 1978. Zooplankton production in the Bering Sea calculated from 1956-1970 Oshoro Maru data. *Marine Science Communications* 4, 329-346.

Irons, D. B. 1992. Aspects of foraging behavior and reproductive biology of the Black-legged Kittiwake. PhD. Dissertation. Univ. of California, Irvine. 143 pp

Iverson, R.L., Coachman, L.K., Cooney, R.T., English, T.S., Goering, J.J., Hunt, G.L., Jr., Macauley, M.C., McRoy, C.P., Reeburgh, W.S., Whitedge, T.E., 1979a.

Ecological significance of fronts in the southeastern Bering Sea. In: Livingston, R.J. (Ed.), *Ecological Processes in Coastal and Marine Systems*. Plenum Press, New York, pp. 437-466.

Iverson, R.L., Whitedge, T.E., Goering, J.J., 1979b. Chlorophyll and nitrate fine structure in the southeastern Bering Sea shelf break front. *Nature* 281, 664-666.

Iverson, S.J., Frost, K.J., Lang, S.L.C. 2002. Fat content and fatty acid composition of forage fish and invertebrates in Prince William Sound, Alaska: factors contributing to among and within species variability. *Marine Ecology Progress Series* 241: 161-181.

Johnson, G.C., Stabeno, P.J., Riser, S.C. Submitted. The Bering Slope Current System Revisited. *Journal of Physical oceanography*.

Kachel, N.B., Hunt, G.L., Jr., Salo, S.A., Schumacher, J.D., Stabeno, P.J., Whitedge, T.E. 2002. Characteristics and variability of the inner front of the southeastern Bering Sea. *Deep Sea Research Part II*. 49: 5889-5909.

Kajimura, H., 1984. Opportunistic feeding of the northern fur seal, *Callorhinus ursinus*, in the eastern North Pacific Ocean and eastern Bering Sea. U.S. Dept. Commerce, NOAA Technical Report NMFS SSRF-779. 49pp.

Khen, G.V., 1987. Interannual changes in water temperatures in the southeastern Bering Sea and its role in oscillation in year class strength of eastern Bering Sea

pollock. Populyationnaya struktura, dinamika chislennosti i ekologiya mintaya [Population structure, numerical dynamics and ecology of pollock]. Vladivostok, pp. 209-220.

Kinder, T.H., Hunt, G.L., Jr., Schneider, D., Schumacher, J.D., 1983. Correlation between seabirds and oceanic fronts around the Pribilof Islands, Alaska. *Estuarine Coastal and Shelf Science* 16, 309-319.

Kinder, T.H., Schumacher, J.D., 1981. Hydrographic structure over the continental shelf of the southeastern Bering Sea. In: Hood, D.W., Calder, J.A. (Eds.), *The Eastern Bering Sea Shelf: Oceanography and Resources*, Vol. 1, US. Government Printing Office, Washington, D.C., pp. 31-52.

Kitaysky, A. S., Piatt, J.F., Wingfield, J.C., and Romano, M. 1999a. The adrenocortical stress-response of black-legged kittiwake chicks in relation to dietary restrictions. *Journal of Comparative Physiology Series B* 169:303-310.

Kitaysky, A. S., Wingfield, J.C., and Piatt, J.F. 1999b. Dynamics of food availability, body condition and physiological stress response in breeding black-legged kittiwakes. *Functional Ecology* 13:577-584.

Kitaysky, A. S., G. L. Hunt, Jr., E. N. Flint, M. A. Rubega, and M. B. Decker. 2000. Resource allocation in breeding seabirds: responses to fluctuations in their food supply. *Marine Ecology Progress Series* 206:283-296.

Kitaysky, A. S., E. V. Kitaiskaia, J. C. Wingfield, and J. F. Piatt. 2001a. Dietary restriction causes chronic elevation of corticosterone and enhances stress response in red-legged kittiwake chicks. *Journal of Comparative Physiology Series B* 171:701-709.

Kitaysky, A. S., J. C. Wingfield, and J. F. Piatt. 2001b. Corticosterone facilitates begging and affect resource allocation in the black-legged kittiwake. *Behavioural Ecology* 12: 619-625.

Kitaysky, A.S., Piatt, J.F., and Wingfield, J.C. 2002. Are seabirds breeding in the southeastern Bering Sea food-limited? Final Report to the North Pacific Marine Research Program.

Kruse, G.H. 1998. Salmon run failures in 1997-98: a link to anomalous conditions? *Alaska Fisheries bul.* 5: 55-63.

Ladd, C., Hunt, G.L., Jr., Stabeno, P.J. Submitted. Climate, mixing, and phytoplankton on the southeastern Bering Sea shelf: A synthesis. Marine Ecology Progress Series

Laevastu, T., Bax, N., 1991. Predation-controlled recruitment in the Bering Sea fish ecosystem. ICES Marine Science Symposium 193, 147-152.

Lang, G.M., Brodeur, R.D., Napp, J.M., Schabetsberger, R., 2000. Variation in groundfish predation on juvenile walleye pollock relative to hydrographic structure near the Pribilof Islands, Alaska. ICES Journal of Marine Science. 57, 265-271.

Leatherwood, S., Bowles, A.E., Reeves, R.R., 1983. Aerial surveys of marine mammals in the southeastern Bering Sea. National Oceanic and Atmospheric Administration/Outer Continental Shelf Environmental Assessment Program, Environmental Assessment of the Alaskan Continental Shelf. Final Reports of Principal Investigators 42, 147-473.

Libes, S.M., 1992. An introduction to marine biogeochemistry. John Wiley and Son, New York, pp.734.

Livingston, P.A., Jurado-Molina, J., 2000. A multispecies virtual population analysis of the eastern Bering Sea. ICES Journal of Marine Science. 57, 294-299.

Livingston, P.A., Lang, G.M., 1996. Interdecadal comparisons of walleye pollock, *Theragra chalcogramma*, cannibalism in the eastern Bering Sea. In: Brodeur, R.D., Livingston, P.A. Loughlin, T.R., Hollowed, A.B. (Eds.), Ecology of juvenile walleye pollock, *Theragra chalcogramma*. NOAA Technical Report NMFS 126, 115-124.

Livingston, P.A., Low, L-L., Marasco, R.J., 1999. Eastern Bering Sea trends. In Sherman, K., Tang, Q. (Eds.), Large Marine Ecosystems of the Pacific Rim:

Assessment, Sustainability, and Management. Blackwell Science, Inc. Malden, Massachusetts, USA, pp. 140-162.

Livingston, P.A., Methot, R., 1998. Incorporation of predation into an assessment model of eastern Bering Sea walleye pollock. In Fishery stock assessment models, Alaska Sea Grant College Program Report No. 98-01, pp.663-678.

Livingston, P.A., Tjelmeland, S., 2000. Fisheries in boreal ecosystems. ICES Journal of Marine Science. 57, 619-627.

Loeng, H., 1989. The influence of temperature on some fish population parameters in the Barents Sea. Journal of Northwest Atlantic Fisheries Science 9, 103-113.

Loughlin, T.R., Antonelis, G.A., Baker, J.D., York, A.E., Fowler, C.W., DeLong, R.L., & Braham, H.W., 1994. Status of the northern fur seal population in the United States during 1992. In: Sinclair, E.H., (Ed.), Fur Seal Investigations, 1994. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-AFSC-45.

Macklin, S.A., 1999. Bering Sea FOCI. In: Loughlin, T.R., Ohtani, K. (Eds.), Dynamics of the Bering Sea. Univ. of Alaska Sea Grant AK-SG-99-03, Fairbanks, Alaska, USA, pp.733-751.

Macklin, S.A., Hunt, G.L., Jr., and Overland, J.E., 2002. Collaborative research on the pelagic ecosystem of the southeastern Bering Sea shelf. Deep Sea Research Part II In Press.

McFarlane, G.A., King, J.R., Beamish, R.J., 2000. Have there been recent changes in climate? Ask the fish. Progress in Oceanography 47, 147-169.

- Mantua, N.J., Hare, S.R., Zhang, Y., Wallace, J.M., Francis, R.C., 1997. A Pacific interdecadal oscillation with impacts on salmon production. *Bulletin American Meteorological Society* 78, 1069-1079.
- Martin, J.E., Grauman, R.D., Marsili, N. 2001. Surface cyclosis in the North Pacific Ocean. Part I: A synoptic climatology. *Mon. Weather Review* 129: 748-765.
- McLaren, I.A., 1963. Effects of temperature on growth of zooplankton and the adaptive significance of vertical migration. *Journal of the Fisheries Research Board of Canada* 20, 685-727.
- McRoy, C.P., Mizobata, K., Nakanishi, S., Saito, S-i., 2001. Trends in ocean conditions and new production on the shelf of the southeastern Bering Sea over the past twenty years. PICES Annual Meeting 2001, Victoria, British Columbia. Abstracts.
- Merrick, R., 1995. The relationship of the foraging ecology of Stellar sea lions (*Eumetopias jubatus*) to their population decline in Alaska. Ph.D. Thesis, Univ. Washington, Seattle, Washington, USA, unpublished.
- Merrick, R.L., 1997. Current and historical roles of apex predators in the Bering Sea ecosystem. *Journal of Northwest Atlantic Fisheries Science* 22, 343-355.
- Merrick, R.L., Chumbley, M.K., Byrd, G.V., 1997. A potential relationship between the diet diversity of Steller sea lions (*Eumetopias jubatus*) and their population decline in Alaska. *Canadian Journal of Fisheries and Aquatic Science* 54, 1342-1348.
- Methven, D.A., Piatt, J., 1991. Seasonal abundance and vertical distribution of capelin (*Mallotus villosus*) in relation to water temperature at a coastal site off eastern Newfoundland. *ICES Journal of Marine Science*. 48, 187-193.

Minobe, S., 1999: Resonance in bidecadal and pentadecadal climate oscillations over the North Pacific: Role in climate regime shifts. *Geophys. Res. Lett.* 26, 855-858.

Mitchell-Innes, B.A., Winter, A. 1987. Coccolithophores: a major component in mature upwelled waters off Cape Peninsula, South Africa in March, 1983. *Marine Biology* 95: 25-30.

Mock, D. W., H. Drummond, C. H. Stinson. 1990. Avian siblicide. *American Scientist* 78:438-449.

Moore, S.E, Waite, M., Mazzuca, L., Hobbs, R.C. 2000. Provisional estimates of Mysticete whale abundance on the central Bering Sea shelf. *Journal of Cetacean Research and Management.* 2: 227-234.

Moore, S.E., Waite, J.M., Friday, N.A., Honkalehto, T. 2002. Cetacean distribution and relative abundance on the central-eastern and the southeastern Bering Sea shelf with reference to oceanographic domains. *Progress in oceanography* 55: 249-261.

NMFS (National Marine Fisheries Service), 1993. Final conservation plan for the northern fur seal (*Callorhinus ursinus*). Prepared by the NMFS/Alaska Fisheries Science Center/National Marine Mammal Laboratory, Seattle, Wash. and the NMFS/Office of Protected Resources, Silver Spring, Md. 80pp.

NMFS (National Marine Fisheries Service). 1995. Status review of the United States Steller sea lion (*Eumetopias jubatus*) population. Prepared by the NMFS/Alaska Fisheries Science Center/National Marine Mammal Laboratory, Seattle, Wash. 45 pp.

NMFS (National Marine Fisheries Service)2002. Fewer northern fur seal pups born in 2002. National Marine Fisheries Service, Alaska Region Press Release, 19 December, 2002, <http://www.fakr.noaa.gov>.

NPFMC (North Pacific Fishery Management Council) 2000. Stock assessment and Fishery Evaluation Document.

NRC (National Research Council), 1996. The Bering Sea ecosystem. National Academy Press, Washington, D.C.

NRC (National Research Council), 2003. decline of the Steller sea lion in Alaskan waters: Untangling food webs and fishing nets. National Academy Press, Washington, D.C.

Nakatani, T., Maeda, T. 1984. Thermal effect on the development of walleye pollock eggs and on their upward speed to the surface. Bulletin Japanese Society of Scientific Fisheries 50: 937-942.

Napp, J. M., Hunt, G. L. Jr., 2001. Anomalous conditions in the south-eastern Bering Sea, 1997: Linkages among climate, weather, ocean, and biology. Fisheries Oceanography 10, 61-68.

Napp, J.M., Incze, L.S., Ortner, P.B., Siefert, D.L. W., Britt, L., 1996. The plankton of Shelikof Strait, Alaska: standing stock, production, mesoscale variability and their relevance to larval fish survival. Fisheries Oceanography 5 (Suppl. 1), 19-38.

Napp, J.M., Kendall, A.W., Jr., Schumacher, J.D., 2000. A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea. Fisheries Oceanography 9, 147-162.

Napp, J.M., Baier, C.T., Coyle, K.O., Brodeur, R.D., Shiga, N., Mier, K., 2002. Interannual and decadal variability in zooplankton communities of the southeastern Bering Sea. Deep-Sea Research Part II. 49: 5991-6008.

Naumenko, N.I., 1996. Distribution, biological condition, and abundance of capelin (*Mallotus villosus*) in the Bering Sea. In: Mathisen, O.A., Coyle, K.O. (Eds.), Ecology of

the Bering Sea: A review of Russian literature. Univ. Alaska Sea Grant Report 96-01, Fairbanks, Alaska, USA, pp. 237-256.

Niebauer, H.J. 1983. Multiyear sea ice variability in the eastern Bering Sea: an update. *Journal of geophysical Research* 88: 2733-2742.

Niebauer, H.J. 1998. Variability in Bering Sea ice cover as affected by a regime shift in the North Pacific in the period 1947-1996. *Journal of Geophysical Research* 103, C12: 27,717-27,737.

Niebauer, H.J., Alexander, V., Cooney, R.T. 1981. Primary production at the eastern Bering Sea ice edge: The physical and biological regimes. Pp. 763-780 In: Hood, D.W., Calder, J.A. (Eds.), *The Eastern Bering Sea Shelf: Oceanography and Resources*, Vol. 1, US. Government Printing Office, Washington, D.C.,

Niebauer, H.J., Alexander, V., Henrichs, S.M., 1990. Physical and biological oceanographic interaction in the spring bloom at the Bering Sea marginal ice edge zone. *Journal of Geophysical Research* 95, 22229-22242.

Niebauer, H.J., Alexander, V., Henrichs, S.M., 1995. A time-series study of the spring bloom at the Bering Sea ice edge. I: Physical processes, chlorophyll and nutrient chemistry. *Continental Shelf Research* 15, 1859-1878.

Niebauer, H.J., Bond, N.A., Yakunin, L.P., Plotnikov, V.V., 1999. An update on the climatology and sea ice of the Bering Sea. In: Loughlin, T.R., Ohtani, K. (Eds.), *Dynamics of the Bering Sea: A Summary of Physical, Chemical, and Biological Characteristics, and a Synopsis of Research on the Bering Sea*. North Pacific Marine Science Organization (PICES), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, USA, pp. 29-59.

Nishiyama, T., 1974. Energy requirement of Bristol Bay sockeye salmon in the central Bering Sea and Bristol Bay In: Hood, D.W., Kelly, E.J. (Eds.), Oceanography of the Bering Sea with emphasis on renewable resources. Institute of Marine Sciences Occasional Publication 2, Univ. Alaska, Fairbanks, Alaska, USA, pp. 321-343.

Ohtani, K., Azumaya, T., 1995. Influence of interannual changes in ocean conditions on the abundance of walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea. In: Beamish, R.J. (Ed.), Climate change and northern fish populations. Canadian Special Publication in Fisheries and Aquatic Science 121: 87-95.

Okkonen, S.R. 2001. Altimeter observations of the Bering Slope Current eddy field. J. geophys. Research 106:2465-2476.

Olson, M.B., Strom, S.L. 2002. Phytoplankton growth, microzooplankton herbivory and community structure in the southeast Bering Sea: insight into the formation and temporal persistence of an *Emiliania huxleyi* bloom. Deep-Sea Research II 49: 5969-5990.

Overland, J.E. and Pease, C.H. 1982. Cyclonic climatology of the Bering Sea and its relation to sea ice extent. Mon. Weather Res. 110: 5-13.

Overland, J.E., Adams, J.M., Bond, N.A., 1999. Decadal variability of the Aleutian low and its relation to high-latitude circulation. Journal of Climate 12, 1542-1548.

Overland, J.E., N.A. Bond, and J.M. Adams, 2001: North Pacific atmosphere and SST anomalies in 1997: Links to ENSO. *Fish. Oceanogr.*, 10, 69-80.

Overland, J.E., N.A. Bond, and J.M. Adams. 2002 : The relation of surface forcing of the bering Sea to large-scale climate patterns. Deep-sea Research II 49: 5855-5868.

Overland, J.E, Spillane, M.C., Hurlburt, H.E., and Wallcraft, A.J. 1994. A numerical study of the circulation of the Bering Sea basin and exchange with the North Pacific Ocean. *Journal of Physical Oceanography* 24: 736-758.

Pahlke, K.A., 1985. Life history and distribution of capelin (*Mallotus villosus*) in Alaskan waters. M.S. Thesis, Univ. Alaska, Juneau, Alaska, USA, Unpublished.

Parker, D.E., Folland, C.K., Ward, M.N., Maskell, K., Bevan, A., Jackson, M., 1993. Ocean-atmosphere climatic fluctuations on interannual to century time scales. Rep. 69, Hadley Centre for Climate Prediction and Research, Bracknell, England, 69 pp.

Pease, C.H., 1980. Eastern Bering Sea ice processes. *Monthly Weather Review* 108, 2015-2023.

Percival, D.B., J.E. Overland and H.O. Mofjeld, 2001: Interpretation of North Pacific variability as a short- and long-memory process. *J. Climate* 14, 4545-4559.

Peterson, W.T., Kister, J.E., Feinberg, L.R. 2002. The effects of the 1997-1999 El Niño- La Niña events on the hydrography and zooplankton off the Central Oregon coast. Prog. In *Oceanography* 54: 381-398.

Piatt, J.F., Anderson, P., 1996. Response of common murrelets to the Exxon Valdez oil spill and long-term changes in the Gulf of Alaska marine ecosystem. *American Fisheries Society Symposium* 18, 720-737.

Quinn, T.J., Niebauer, H.J., 1995. Relation of eastern Bering Sea walleye pollock (*Theragra chalcogramma*) recruitment to environmental and oceanographic variables. In: Beamish, R.J. (Ed.), *Climate change and northern fish populations*. Canadian Special Publication in Fisheries and Aquatic Science 121, 497-507.

Reed, R.K., 1995. On the variable subsurface environment of fish stocks in the Bering Sea. *Fisheries Oceanography* 4, 317-323.

Reed, R.K., 1998. Confirmation of a convoluted flow over the Southeastern Bering Sea Shelf. *Continental Shelf Research* 18, 99-103.

Reed, R.K., In Press???. Confirmation of a convoluted flow over the southeastern Bering Sea shelf. *Continental Shelf Research* ???

Reed, R.K. and Stabeno, P.J., 1996. On the climatological mean circulation over the eastern Bering Sea shelf. *Continental Shelf Research* 16, 1297-1305.

Reed, R.K. and Stabeno, P.J., 1999. The Aleutian North Slope Current. In Loughlin, T.R. and Ohtani, K. (eds). *Dynamics of the Bering Sea*. University of Alaska Sea Grant, AK-SG-99-03, pp. 177-192.

Reid, P.C., Battle, E.J.V., Batten, S.D., Brander, K.M., 2000. Impacts of fisheries on plankton community structure. *ICES Journal of Marine Science* 57, 495-502.

Rho, T., 2000. Carbon and nitrogen uptake dynamics during 1997 and 1998: Anomalous conditions in the Bering Sea. M.S. Thesis, University of Alaska Fairbanks. 95 pp.

Richardson, K., Christoffersen, A., 1991. Seasonal distribution and production of phytoplankton in the southern Kattegat. *Marine Ecology Progress Series* 78, 217-227.

Rivkin, R.B., Putland, J.N., Anderson, M.R., Deibel, D. 1999. Microzooplankton bacterivory and herbivory in the NE subarctic Pacific. *Deep Sea Research II* 46: 2579-2618.

Roach, A.T., Aagaard, K., pease, C.H., Salo, S.A., Weingartner, T., Pavlov, V., Kulakov, M. 1995. Direct measurements of transport and water properties through Bering Strait. *J. Geophysical research* 100: 18,443-18,457.

Robertson, J.E., Robinson, C., Turner, D.R., Holligan, P., Watson, A.J., Boyd, P., Fernandez, E. and Finch, M. (1994) The impact of a coccolithophore bloom on oceanic carbon uptake in the northeast Atlantic during summer 1991. *Deep-Sea Res.* **41**:297-314.

Robson, B.W., 2002. The relationship between foraging areas and breeding sites of lactating northern fur seals, *Callorhinus ursinus* in the eastern Bering Sea. MSc. Thesis, University of Washington, Seattle, WA.

Roff, J.C., Middlebrook, K., Evans, F., 1988. Long-term variations in North Sea zooplankton. *Journal of the Marine Biology Association of the U.K.* 68, 143-164.

Runge, J.A., 1988. Should we expect a relationship between primary production, fisheries? The role of copepod dynamics as a filter of trophic variability. *Hydrobiologia* 167/168, 61-71.

Russell, R.W., Harrison, N.M., and Hunt, G.L., Jr. 1999. Foraging at a front: hydrography, zooplankton and avian planktivory in the northern Bering Sea. *Mar. Ecol. Prog. Ser.* 182: 77-93.

Sætersdal, G., Loeng, H., 1987. Ecological adaptation of reproduction in northeast Arctic cod. *Fisheries Research* 5, 253-270.

Sambroto, R.N., Niebauer, H.J., Goering, J.J., Iverson, R.L., 1986. Relationships among vertical mixing, nitrate uptake and phytoplankton growth during the spring bloom in the southeast Bering Sea middle shelf. *Continental Shelf Research* 5, 161-198.

Schabetsberger, R., Brodeur, R.D., Ciannelli, L., Napp, J.M., Swartzman, G.L., 2000. Diel vertical migration and interactions of zooplankton and juvenile walleye pollock (*Theragra chalcogramma*) at a frontal region near the Pribilof Islands, Bering Sea. ICES Journal of Marine Science 57, 1283-1295.

Schell, D.M., 2000. Declining carrying capacity in the Bering Sea: Isotopic evidence from whale baleen. Limnology and Oceanography 45, 459-462.

Schell, D.M., 2001. Carbon isotope ratio variations in the Bering Sea biota: The role of anthropogenic carbon dioxide. Limnology and Oceanography 46: 999-1000.

Schell, D.M., Barnett, B.A., Vinette, K. 1998. Carbon and nitrogen isotope ratios in zooplankton of the Bering, Chukchi, and Beaufort Seas. Marine Ecology Progress Series 162: 11-23.

Sherr, E.B., Sherr, B.F. 1992. Trophic roles of pelagic protists: phagotrophic flagellates as herbivores. Arch. Hydrobiol. Beih Ergebn Limnol 37: 165-172.

Schumacher, J.D., and Alexander, V. 1999. Variability and the role of the physical environment in the Bering Sea ecosystem. In: Loughlin, T.R., Ohtani, K. (Eds.), Dynamics of the Bering Sea: A Summary of Physical, Chemical, and Biological Characteristics, and a Synopsis of Research on the Bering Sea, North Pacific Marine Science Organization (PICES), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, USA, pp. 147–160.

Schumacher, J.D., Reed, R.K. 1992. Characteristics of currents over the continental slope of the eastern Bering Sea. J. Geophys. Oceanogr. 97: 9423-9433.

Schumacher, J.D., and Stabeno, P.J., 1994. Ubiquitous eddies of the eastern Bering Sea and their coincidence with concentrations of larval pollock. *Fisheries Oceanography* 3, 182-190.

Schumacher, J., Stabeno, P.J., 1998. The continental shelf of the Bering Sea, In: *The Sea*, Vol. XI, The Global Coastal Ocean: Regional Studies, Synthesis, J. Wiley, New York, New York, USA, pp.789-823.

Schumacher, J.D., Bond, N.A., Brodeur, R.D., Livingston, P.A., Napp, J.M., Stabeno, P.J., In Press. Climate change in the southeastern Bering Sea and some consequences for biota. In: Hemple, G., Sherman, K. (Eds.), *Large marine ecosystems of the world: trends in exploitation, protection and research*. Amsterdam: Elsevier Science.

Schumacher, J.D., Kinder, T.H., Pashinski, D.J., Charnell, R.L., 1979. A structural front over the continental shelf of the eastern Bering Sea. *Journal of Physical Oceanography* 9, 79-87.

Schwing, F.B. and Moore, C.S. 2000. A year without summer for California: or a harbinger of a climate shift. *Transactions of the American Geophysical Union* 81: 301.

Shuert, P.G. and Walsh, J.J. 1993. A coupled physical-biological model of the Bering-Chukchi Seas. *Continental Shelf research* 13: 543-574.

Sinclair, E.H., 1988. Feeding habits of northern fur seals in the eastern Bering Sea. M.S. Thesis, College of Oceanography, Oregon State Univ., Corvallis, Oregon, USA, Unpublished.

Sinclair, E.H. and Stabeno, P.J. 2002. Mesopelagic nekton and associated physics of the southeastern Bering Sea. *Deep-Sea Research II* 49: 6127-6145.

Sinclair, E.H., Antonelis, G.A., Robson, B.W., Ream, R.R., Loughlin, T.R., 1996. Northern fur seal, *Callorhinus ursinus*, predation on juvenile walleye pollock, *Theragra chalcogramma*. In: Brodeur, R.D., Livingston, P.A. Loughlin, T.R., Hollwed, A.B. (Eds.), Ecology of juvenile walleye pollock, *Theragra chalcogramma*. NOAA Technical Report NMFS 126, 167-178.

Sinclair, E.H., Balanov, A.A., Kubodera, T., Radchenko, V.I., Fedorets, Y.A. 1999. Distribution and ecology of mesopelagic fishes and cephalopods. In: . Loughlin, T.R., Ohtani, K. (Eds.), Dynamics of the Bering Sea: A Summary of Physical, Chemical, and Biological Characteristics, and a Synopsis of Research on the Bering Sea, North Pacific Marine Science Organization (PICES), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, USA, pp. 485-508.

Sinclair, E.H., Loughlin, T., Pearcy, W., 1994. Prey selection by northern fur seals (*Callorhinus ursinus*) in the eastern Bering Sea. Fisheries Bulletin 92, 144-156.

Smith, S.L., Vidal, J., 1984. Spatial and temporal effects of salinity, temperature, and chlorophyll on the communities of zooplankton in the southeastern Bering Sea. Journal of Marine Research 42, 221-25784:

Smith, S.L., Vidal, J., 1986. Variations in the distribution, abundance and development of copepods in the southeast Bering Sea in 1980 and 1981. Continental Shelf Research 5, 215-239.

Smith, S.L., Henrichs, S.M., Rho, T. 2002. Stable C and N isotopic composition of sinking particles and zooplankton over the southeastern Bering Sea shelf. Deep-Sea research II 49: 6031-6050.

Sogard, S.M., Olla, B.L. 1993. Effects of light, thermoclines and predator presence on vertical distribution and behavioral interactions of juvenile walleye pollock, *Theragra chalcogramma* Pallas. J. Experimental Marine Biology and Ecology 167: 179-195.

Spencer, P.D., Collie, J.S., 1996. A simple predator-prey model of exploited marine fish populations incorporating alternative prey. ICES Journal of Marine Science 53, 615-628.

Springer, A.M., 1992. A review: Walleye pollock in the North Pacific- how much difference do they really make? Fisheries Oceanography 1, 80-96.

Springer, A.M., 1998. Is it all climate change? Why marine bird and mammal populations fluctuate in the North Pacific. In: Holloway, G., Muller, P., Henderson, D. (Eds.), Proceedings of the 10<sup>th</sup> 'Aha Huliko'a Hawaiian Winter Workshop on Biotic Impacts of Extratropical Climate Variability in the Pacific, January 26-30, 1998. SOEST Special Publication, pp.121-125.

Springer, A.M., McRoy, C.P., Flint, M.V., 1996a. The Bering Sea green belt: shelf-edge processes and ecosystem production. Fisheries Oceanography 5, 205-223.

Springer, A.M., Murphy, E.C., Roseneau, D.G., McRoy, C.P., Cooper, B.A. 1987. The paradox of food webs in the northern Bering Sea-I. Seabird food habits. Continental Shelf Research 7(8), 895-911.

Springer, A.M., McRoy, C.P., and Turco, K.R. 1989. The paradox of pelagic food webs in the northern Bering Sea-II. Zooplankton communities. Continental Shelf Research 9, 359-386.

Springer, A.M. and Roseneau, D.G. 1985. Copepod-based food webs: auklets and oceanography in the Bering Sea. Marine Ecology Progress Series 21: 229-237.

Stabeno, P.J., Hunt, G.L., Jr. 2002. Overview of progress realized in the Inner Front and Southeast Bering Sea Carrying Capacity Programs. *Deep Sea Research Part II*. 49: 61-57-6168.

Stabeno, P.J., Overland, J.E. 2001. Bering Sea shifts toward an earlier spring transition. *Eos, Transactions, amer. Geophysical Union* 82: 317, 321.

Stabeno, P.J., Van Meurs, P. 1999. Evidence of episodic on-shelf flow in the southeastern Bering Sea. *J. Geophys. Research* 104 (C12): 29715-29720.

Stabeno, P.J. and Reed, R.K. 1994. Circulation in the Bering Sea basin observed by satellite-tracked drifters: 1986-1993. *Journal of Physical Oceanography* 24: 848-854.

Stabeno, P.J., Kachel, N.B., Sullivan, M., Whitedge, T.E., This Volume. Variability along the 70 m isobath of the southeastern Bering Sea. *Deep Sea Research Part II*

Stabeno, P.J., Schumacher, J.D., Davis, R.F., Napp, J.M., 1998. Under-ice observations of water column temperature, salinity, and spring phytoplankton dynamics: Eastern Bering Sea shelf. *Journal of Marine Research* 56, 239-255.

Stabeno, P.J., Schumacher, J.D., Ohtani, K., 1999a. The physical oceanography of the Bering Sea. In: Loughlin, T.R., Ohtani, K. (Eds.), *Dynamics of the Bering Sea: A Summary of Physical, Chemical, and Biological Characteristics, and a Synopsis of Research on the Bering Sea*, T.R. Loughlin and K. Ohtani (eds.), North Pacific Marine Science Organization (PICES), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, USA, pp. 1–28.

Stabeno, P.J., Schumacher, J.D., Salo, S.A., Hunt, G.L., Jr., Flint, M. (1999b). Physical environment around the Pribilof Islands. Pp. 193-215 In, Loughlin, T. R., and K. Ohtani

(eds.), Alaska Sea Grant Press (Eds.), *The Bering Sea: Physical, Chemical and Biological Dynamics*. University of Alaska Sea Grant, Fairbanks.

Stabeno, P.J., Bond, N.A., Kachel, N.B., Salo, S.A., Schumacher, J.D., 2001. On the temporal variability of the physical environment over the southeastern Bering Sea. *Fisheries Oceanography* 10, 81-98.

Steele, J.H., Henderson, E.W., 1984. Modeling long-term fluctuations in fish stocks. *Science* 224, 985-987.

Stockwell, D. A., Whitley, T.E., Zeeman, S.I., Coyle, K.O., Napp, J.M., Brodeur, R.D., Pinchuk, A.I., Hunt, G.L., Jr., 2001. Anomalous conditions in the south-eastern Bering Sea, 1997: Nutrients, phytoplankton, and zooplankton. *Fisheries Oceanography* 10, 99-116.

Sugimoto, T., Tadokoro, K., 1997. Interannual-interdecadal variations in zooplankton biomass, chlorophyll concentration and physical environment in the subarctic Pacific and Bering Sea. *Fisheries Oceanography* 6, 74-93.

Sugimoto, T., Tadokoro, K., 1998. Interdecadal variations of plankton biomass and physical environment in the North Pacific. *Fisheries Oceanography* 7, 289-299.

Sukhanova, I.N. and Flint, M.V. 1998. Anomalous blooming of coccolithophorids over the eastern Bering Sea shelf. *Oceanography* 38: 502-505.

Sukhanova, I.N., Semina, H.J., Venttsel, M.V. 1999. Spatial distribution and temporal variability of phytoplankton in the Bering Sea. In: Loughlin, T.R., Ohtani, K., (Eds.). *Dynamics of the Bering Sea*. University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, pp. 193-215.

Swartzman, G., Napp, J., Brodeur, R., Winter, A., & Ciannelli, L. (In Press). Spatial patterns of pollock and zooplankton distribution in the Pribilof Islands, Alaska, nursery area and their relationship to pollock recruitment. ICES Journal of Marine Science.

Thompson, D.W.J., Wallace, J.M. 1998. The Arctic Oscillation signature in the wintertime geopotential height and temperature fields. Geophys. Res. Letters 25: 1297-1300.

Tomasada, A., Odate, K., 1995. Long-term variability in zooplankton biomass and environment. *Umi to Sora* 71, 1-7. (in Japanese)

Tortell, P.D., Rau, G.H., Morel, F.M.M., 2000. Inorganic carbon acquisition in coastal Pacific phytoplankton communities. Limnology and Oceanography 45, 1485-1500.

Townsend, D.W., Cammen, L.M., Holligan, P.M., Campbell, D.E., Pettigrew, N.R., 1994. Causes and consequences of variability in the timing of spring phytoplankton blooms. Deep-Sea Research 41, 747-765.

Townsend, T.W., Keller, M.D., Holligan, P.M., Ackleson, S.G., Balch, W.M. 1994. Blooms of the coccolithophore *Emiliana huxleyi* with respect to hydrography in the Gulf of Maine. Cont. Shelf Res. 14: 979-1000.

Trenberth, K.E., Hurrell, J.W., 1995. Decadal coupled atmospheric-ocean variations in the North Pacific Ocean. In: Beamish, R.J. (Ed.), Climate change and northern fish populations. Canadian Special Publication in Fisheries and Aquatic Science 121, 15-24.

Trites A. W., Livingston P. A., Vasconcellos M. C., Mackinson S., Springer A. M., Pauly D., 1999. Ecosystem change and the decline of marine mammals in the Eastern Bering Sea: testing the ecosystem shift and commercial whaling hypotheses. Fisheries Centre Reports 7 (1), 98 pp.

Turnock, B. J., Wilderbuer, T. K., Brown, E.R., 2000. Arrowtooth flounder. In: Stock Assessment and Fishery Evaluation (SAFES) for the Gulf of Alaska and Bering Sea Aleutian Islands, Appendix B, Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska. North Pacific Fishery Management Council, Anchorage, Alaska, USA. Pp. 138-167.

Tynan, C.T. 1998. Coherence between whale distributions, chlorophyll concentration, and oceanographic conditions on the southeast Bering Sea shelf during a coccolithophore bloom, July-August, 1997. EOS 79: 127.

Tynan, C. 1999. Redistribution of cetaceans in the southeast Bering Sea relative to anomalous oceanographic conditions during the 1997 El Nino. In proceedings of the 1998 Science board symposium on the impacts of the 1997.98 El nino event on the North Pacific Ocean and its marginal seas. (Eds. Freeland, H.J., Peterson, W.T., Tyler, A.) PICES Scientific Report 10: 115-117.

Van Pelt, T.I., Piatt, J.F., Lance, B.K., & Roby, J.F. 1997. Proximate composition and energy density of some North Pacific forage fish. Comparative Biochemistry and Physiology 118A, 1393-1398.

Vance, T.C., Baier, C.T., Brodeur, R.D., Coyle, K.O., M. B. Decker, M.B., Hunt, G.L., Jr., Napp, J.M., Schumacher, J.D., Stabeno, P.J., Stockwell, D., Tenant, C.T., Whitley, T.E., Wyllie-Echeverria, T., Zeeman, S., 1998. Anomalies in the Ecosystem of the Eastern Bering Sea: Including Blooms, Birds and Other Biota. EOS, Transactions of the American Geophysical Union 79 (10), 121/126.

Vidal, J., 1980. Physioecology of zooplankton II: Effects of phytoplankton concentration, temperature, and body size on the development and molting rates of *Calanus pacificus* and *Pseudocalanus* spp. Marine Biology 56, 135-146.

Vidal, H., Smith, S.L., 1986. Biomass, growth and development of population of herbivorous zooplankton in the southeastern Bering Sea. *Deep-Sea Research* 33, 523-566.

Vlietstra, L., Springer, A., Byrd, G.V., Hunt, G.L., Jr. 2003. Control of seabird population dynamics at the Pribilof Islands. Final Report for the seabird section of the Synthesis of information on Marine birds and Fur Seals and Steller sea lions in the eastern Bering Sea. SEBSCC report. **THIS will be formalized as a “publication”**

Wallace, J.M., C. Smith and C.S. Bretherton, 1992: Singular value decomposition of wintertime sea surface temperature and 500-mb height anomalies. *J. Climate* 5, 561-576.

Walsh, J.J., McRoy, C.P., 1986. Ecosystem analysis in the southeastern Bering Sea. *Continental Shelf Research* 5, 259-288.

Wespestad, V.G., 1991. Pacific herring population dynamics, early life history and recruitment variation relative to eastern Bering Sea oceanographic factors. Ph.D. Thesis, Univ. of Washington, Seattle, Washington, USA, Unpublished.

Wespestad, V.G., 1994. Walleye Pollock. Report and data submitted for the Third International Pollock Assessment Workshop. Seattle, WA. Pp. 119-143.

Wespestad, V.G., Fritz, L.W., Ingraham, W.J., Megrey, B.A., 2000. On relationships between cannibalism, climate variability, physical transport, and recruitment success of Bering sea walleye pollock (*Theragra chalcogramma*). *ICES Journal of Marine Science*. 57, 272-278.

Whitledge, T.E., 1999. Nutrient cycling and phytoplankton growth. In: Macklin, S.A. (Ed.), Report on the FOCI International Workshop on Recent Conditions in the Bering Sea, Pacific Marine Environmental Laboratory, Seattle, pp. 13-14.

Whitledge, T.E., Reeburgh, W.S., Walsh, J.J., 1986. Seasonal inorganic nitrogen distribution and dynamics in the southeastern Bering Sea. *Continental Shelf Research* 5, 109-132.

Wilderbuer, T.K., Hollowed, A.B., Ingraham, W.J., Jr., Spencer, P.D., Connors, M.E., Bond, N.A., Walters, G.E. 2002. Flatfish recruitment response to decadal climate variability and ocean conditions in the eastern Bering Sea. *Progress in Oceanography* 55: 235-247.

Witherell, D., Pautzke, C., Fluharty, D., 2000. An ecosystem-based approach for Alaska groundfish fisheries. *ICES Journal of Marine Science*. 57, 771-777.

Withrow, D.E., Loughlin, T.R., 1995. Abundance and distribution of harbor seals (*Phoca vitulina richardsi*) along the north side of the Alaska Peninsula and Bristol Bay during 1995. In: Marine Mammal Protection Act and Endangered Species Act Implementation Program, 1995. National Marine Mammal Laboratory, National Marine Fisheries Service/NOAA, Seattle WA, November 1996.

Wyllie-Echeverria, T., 1995. Sea-ice conditions and the distribution of walleye pollock (*Theragra chalcogramma*) on the Bering and Chukchi shelf. In: Beamish, R.J. (Ed.), Climate change and northern fish populations. Canadian Special Publication in Fisheries and Aquatic Science 121: 131-136.

Wyllie-Echeverria, T., 1996. The relationship between the distribution of one-year-old walleye pollock, *Theragra chalcogramma*, and sea-ice characteristics In: Brodeur, R.D.,

Livingston, P.A. Loughlin, T.R., Hollowed, A.B. (Eds.), Ecology of juvenile walleye pollock, *Theragra chalcogramma*. NOAA Technical Report NMFS 126, 47-56.

Wyllie-Echeverria, T. and Wooster, W.S., 1998. Year-to-year variations in Bering Sea ice cover and some consequences for fish distributions. Fisheries Oceanography 7, 159-170.

York, A.E., Hartley, J.R., 1981. Pup production following harvest of female northern fur seals. Canadian Journal of Fisheries and Aquatic Science 38, 84-90.

York, A.E., Kozloff, P., 1986. On estimating the number of fur seal pups born on St. Paul Island 1980-1986. Fishery bulletin 85, 367-375.

York, A.E., Towell, R.G., Ream, R.R., Baker, J.D., Robson, B.W., 2000. Population assessment, Pribilof Islands, Alaska. In: Robson, B.W. (Ed.) Fur Seal Investigations, 1998. U.S. Department Commerce, NOAA Technical Memorandum NMFS-AFSC-113, pp. 7-26.

Table 1. Relationship between the timing of ice retreat and the type of spring bloom. From hunt and Stabeno 2002b.

	Bloom Occurs at the Ice Edge	Bloom in Open Water
Ice Gone by mid-March	0	7
Ice remains after late March	6	0

Table 2. Comparison of large-scale *Emiliania huxleyi* blooms. (After Napp and Hunt, 2001)

<b><u>Size (km<sup>2</sup>)</u></b>		<b><u>Source</u></b>
Coastal Blooms	5-10 x 10 <sup>4</sup>	Holligan <i>et al.</i> , 1983
	5 x 10 <sup>5</sup>	Balch <i>et al.</i> , 1991
	5 x 10 <sup>5</sup>	Brown & Yoder, 1993
	5 x 10 <sup>5</sup>	Brown & Podesta, 1997
Open Ocean Blooms Bering Sea 1997	5 x 10 <sup>5</sup>	Holligan <i>et al.</i> , 1993
	2.4 x 10 <sup>5</sup>	Napp and Hunt, 2001
<b><u>Duration of Bloom (mo.)</u></b>		
Gulf of Maine, 1998	□ 1	Townsend <i>et al.</i> , 1994
Gulf of Maine, 1989	□ 1	Townsend <i>et al.</i> , 1994
NE Atlantic, 1991	1.1	Robertson <i>et al.</i> , 1994
Bering Sea, 1997	4	Napp and Hunt, 2001
<b><u>Cell Density (cells ml<sup>-1</sup>)</u></b>		
Gulf of Maine, 1988	2.4 x 10 <sup>3</sup>	Townsend <i>et al.</i> , 1994
Gulf of Maine, 1989	1.5 x 10 <sup>3</sup>	Townsend <i>et al.</i> , 1994
NE Atlantic, 1991	1.0 x 10 <sup>3</sup>	Robertson <i>et al.</i> , 1994
Bering Sea, 1997	3.1-4.7 x 10 <sup>3</sup>	Napp and Hunt, 2001
<b><u>Lith Density (liths ml<sup>-1</sup>)</u></b>		
Gulf of Maine, 1988	1.3 x 10 <sup>5</sup>	Townsend <i>et al.</i> , 1994
Gulf of Maine, 1989	3.0 x 10 <sup>5</sup>	Townsend <i>et al.</i> , 1994
NE Atlantic, 1991	3.5 x 10 <sup>5</sup>	Robertson <i>et al.</i> , 1994
Bering Sea, 1997	3 - 5 x 10 <sup>5</sup>	Napp and Hunt, 2001

Table 3. Responses of calanoid copepods to interannual variation in water temperature during the spring bloom in the Bering Sea. Copepod data are numbers m<sup>-3</sup> from the middle shelf in May 1980 and 1981. \* = difference significant p < 0.05. Data from Smith and Vidal (1986).

Variable	Year		difference
	1980	1981	
Onset of Bloom	25 April	5 May	
Termination of Bloom	28 May	29 May	
Temperature (°C) Prior to Bloom – top 20 m	0.97	3.04	2.07
Temperature During Bloom – top 20 m	2.22	5.06	2.84
<i>Acartia</i> spp. All copepodids	18.9	8.5	- 55%
<i>Acartia</i> spp. Adult males and female	9.4	2.8*	- 70%
<i>Pseudocalanus</i> spp. All copepodids	83.1	308.5*	+ 270%
<i>Calanus marshallae</i> All copepodids	31.7	30.6*	- 3.5%
<i>Calanus marshallae</i> Adult females	0.1	0.8*	+ 700%
<i>Metridea pacifica</i> all copepodids	1.6	20.3*	+ 1169%
<i>Oithona</i> spp.	269.6	233.4	- 13.4%

Table 4. Responses of calanoid copepods to interannual variation in water temperature during the spring bloom in the Bering Sea. Copepod data are numbers m<sup>-3</sup> from the middle shelf and inner shelf in June 1997, 1998 and 1999. \* = difference significant p < 0.05. Data from Coyle and Pinchuk (2002b) and Hunt et al. (2002a).

Variable	Year			Change in 1999 from mean of 1997 + 1998
	1997	1998	1998	
Onset of Bloom	Mid - April	Early May	Late March	
Temperature During June, integrated water column °C	3.76	3.45	0.32*	3.29
<i>Acartia</i> spp.	961	711	64*	- 92%
<i>Pseudocalanus</i> spp.	1168	893	240*	- 77%
<i>Calanus marshallae</i>	34	72	3.7*	- 93%
<i>Calanoid nauplii</i>	616	626	322*	- 48%
<i>Oithona similis</i>	99	219*	28	- 82%
Chaetognaths	43	12*	28	+ 1.8%

Table 5. Comparison of acoustically determined euphausiid biomass among the years 1997, 1998 and 1999, mean and 95% confidence interval. \* = Statistically significant at  $p < 0.05$ . Data from Coyle and Pinchuk (2002a).

	June 1997	June 1998	June 1999
Transect Location	Mean (g.m <sup>-2</sup> ) (95% CI)	Mean (g.m <sup>-2</sup> ) (95% CI)	Mean (g.m <sup>-2</sup> ) (95% CI)
Port Moller, Line A	2.3 (1.94 – 2.57)	0.77 (0.54 – 0.99)	17.27* (13.27 – 21.27)
Port Moller, Line C	2.98 (2.36 – 3.61)	0.34 (0.25 – 0.44)	15.76* (14.60 – 17.46)
Port Moller, Line E	No Data	0.73 (0.62 – 0.83)	5.29* (4.30 – 5.65)
Cape Newenham, Line C	1.91 (1.14 – 1.70)	5.50* (4.71 – 6.22)	2.75 (1.59 – 3.91)
Nunivak Island, Line C	1.58 (1.46 – 1.70)	2.38 (2.26 – 2.49)	6.82* (5.91 – 7.72)
Nunivak Island, Line E	1.25 (1.13 – 1.36)	1.93 (1.62 – 2.19)	5.64* (4.89 – 6.02)

Table 6. Strong year classes ( $\geq 20\%$  above the mean from 1975 to 1999) and weak year-classes ( $\leq 20\%$  below the mean) by species in the Bering Sea. Years not shown had average year classes (mean  $\pm 20\%$ ). Data from NPFMC, 2000.

Group	Strong Year-classes	Weak Year-classes
Group A		
Walleye Pollock	77, 78, 79, 80, 84, 89, 92, 96	75, 76, 81, 83, 85, 87, 88, 91, 93, 94, 98, 99
Alaska Plaice	75, 76, 77, 78, 79, 80, 82, 88, 90	83, 84, 85, 86, 87, 94, 95, 96, 97
Yellowfin Sole	75, 76, 79, 81, 83, 91	78, 82, 84, 85, 86, 89, 90, 94
Flathead Sole	77, 79, 80, 81, 84, 85, 87	75, 76, 88, 90, 91, 92, 93, 94, 95, 96, 97
Pacific Cod	77, 78, 79, 82, 84, 89, 92	81, 83, 86, 87, 88, 91, 93, 94, 95, 97, 98, 99
Group B		
Rock Sole	80, 81, 83, 84, 85, 86, 87, 88, 90	75, 76, 77, 78, 79, 91, 92, 94, 95
Arrowtooth Flounder	80, 81, 83, 84, 86, 87, 88, 91	75, 76, 78, 79, 82, 93, 94, 96, 97
Greenland Turbot	75, 76, 77, 78, 79	81, 82, 83, 86, 87, 88, 89, 91, 92

Table 7. Hypotheses for explaining variation in year-class strength of walleye pollock in the eastern Bering Sea.

Hypothesis	Citations	Tests
<b>A. Production-Based Hypotheses</b>		
1. Decreases in wind mixing negatively influence post-bloom primary production	Sambrotto et al. 1986 Hunt et al., 2002a Springer, This Report, Part 4	Compare wind mixing in years with strong and weak year-classes; Ladd et al., ms. Probably no significant relationship with summer mixing.  Determine if there is strong coupling between spring and summer primary production, zooplankton abundance, and pollock year-class strength.
2. Late ice retreat results in cold water blooms and reduced zooplankton production	Stabeno et al., 1998 Hunt et al., 2002a Hunt and Stabeno, 2002	Compare zooplankton abundances in years with early and late ice retreat; Coyle and Pinchuk 2002b show significant differences  Determine if zooplankton abundance limits age-0 or age-1 pollock survival
<b>B. Cannibalism-based Hypotheses</b>		
3. When the cold pool is large, age-1 pollock move to the shelf edge where adults abound	Dwyer et al., 1987 Ohtani and Azumaya, 1995 Wyllie-Echeverria, 1995	Compare cold pool size in years with strong and weak year classes;
4. Favourable winds in early spring may transport eggs, larvae and small age-0 pollock deep onto the shelf thus separating them from adults	Wespestad et al., 2000	Compare wind forcing in years with strong and weak year classes
<b>C. Mixed production/cannibalism</b>		
<u>Hypothesis</u> 5. Adult pollock switch prey types such that when zooplankton are scarce in spring, they increase cannibalism	Cooney et al., 2001	Since zooplankton are less abundant in cold years, compare the amount of cannibalism in years with cold and warm spring temperatures
<b>D. Mixing Process hypothesis</b>		
6. Intermediate levels of turbulence are required for optimal foraging conditions	Megrey and Hinckley, 2001	Compare turbulence levels in years with strong and weak year-classes




Figure 1. Three dimensional view of the eastern Bering Sea, with location of the biophysical mooring, M2 shown. Map from ?????




Figure 2. Southeastern Bering Sea showing isobathes and domains. Map courtesy of N. Kachel.

Need to add Labels: Unimak Pass, Nunivak island etc GET MAP FROM PHYLLIS THAT SHE USED IN CHAPTER ON BERING SEA




Figure 3. Schematic of major currents in the Bering Sea. From Stabeno et al. 1999a.

Map of currents

**Get rid of W in Bristol bay; add some eddies in the BSC**




Figure 4. Time series of the Arctic Oscillation (AO) and the Pacific Decadal Oscillation (PDO). Note the change from negative to positive in the PDO as of 1977 and from positive to negative in 1998. The AO showed a shift from negative to positive about 1989. From Overland et al. ???




Figure 5. The location of the Aleutian Low, the statistical mean location of winter low pressure cells in the northern North Pacific. From????  
Overland Fig 1




Figure 6. The strength of the Aleutian Low with respect to the long-term mean, annual signal and 5-year running mean. Note that the signal is dominated by extremely strong interannual variability. From Percival et al., 2001 Overland Fig 2




Figure 7. Auto-correlation functions (stick plots) of the Aleutian Low time series, with three models of the time series: left- Auto-Regressive; middle- Long memory; and right- 50-year oscillation plus white noise (From Percival et al., 2001).




Figure 8. Time series of Top- Mean Sea Surface Temperature anomaly; Middle- Downward Shortwave Radiation Anomaly and Bottom- ETC From Bond fig 2

Need proper caption maybe better figure




Figure 9 From Bond **Need to get proper Caption Bond fig 1**




Figure 10. Percent Ice cover for two latitudinal bands in the eastern Bering Sea. Note the decrease in ice cover in the southern region as of about 1977, whereas there is little evidence for a change in ice cover at this time in the more northerly region. From Hunt et al., 2002a.




Figure 11. Number of weeks with ice cover of . Note....from Hunt et al., 2002a.




Figure 12. Time series of ocean temperatures and fluorescence from Mooring 2. The thin yellow line at the bottom of each panel is fluorescence scaled to the maximum each year. Temperature of  $<-1$  °C indicate the presence of melting ice. From Hunt et al., 2002a.




Figure 13. The relationship between the timing of the last winter storm, ice retreat, and the relationship of the bloom to sea ice. Note that blooms that occur to the right of the dotted gray line are associated with the ice edge. From Hunt et al., 2002a.




Figure 14. relationship between timing of ice retreat, whether the bloom will occur in association with ice in cold water or in open, warmer water, and the potential effect of a warm water bloom on copepod production. After Hunt et al., 2002a.






Figure 15. Wind speed cubed at St. Paul island. Note that since the late 1970s winter winds have been less strong and in summer, there have been more periods of wind below the long-term average (those areas in light gray). From Hunt et al., 2002a

**September 1997**



**25 April 1998**



**12 September 1999**




Figure 16. Three SeaWiFS views of the Bering Sea coccolithophore bloom.




Figure 17. SeaWiFS false color image of the coccolithophore bloom, 24 July 1998. Note the trace of coccoliths in the water passing through Shpanberg Strait to the east of St. Lawrence island, and thence northward through the east side of Bering Strait. From Napp and Hunt 2001.




Figure 18. Changes in zooplankton biomass in the deep basin and in the outer, middle and coastal domains of the southeastern Bering Sea sampled by the Oshoro Maru Summer Cruises. Data from 1997 to 1994 from Sugimoto and Tadokoro (1998). Data from 1995 to 1999 from Dr. N. Shiga, unpublished. Means with standard errors. From Hunt et al. (2002a).





Figure 19. June abundances of *Pseudocalanus* spp. (top) and *Acartia* spp. (bottom) in relation to the integrated water temperatures at the stations where they were collected 1997-1999). Dashed lines are 95% confidence intervals around the regressions. K. Coyle, unpublished data.




Figure 20. Biomass of large medusae in the National Marine Fisheries Service bottom trawl surveys of the southeastern Bering Sea. Data courtesy of G. Walters and R. Brodeur.










Figure 21. Time series of bottom trawl surveys in three regions of the southeastern Bering Sea that have been consistently surveyed with similar gear since the mid-1960s. From Connors et al., 2002.















Figure 22. Spawner biomass of selected ground fish in the eastern Bering Sea. Data from NPFMC SAFE 2000.












Figure 23. year-class strength for selected species of groundfish in the eastern Bering Sea. Data from NPFMC SAFE 2000.

# WESTERN ALASKAN SALMON CATCH



from:  
Gordon Kruse  
Alaska Dept. Fish and Game

Figure 24. Time-series for three western Alaska salmon runs. From Kruse, 1998.





Figure 25. Time-series of pup counts at St Paul Island and St. George Island, Pribilof Islands. From NMFS, 2002.




Figure 26. Changes in the number of adult (left scale, filled dots) and pup (right scale, open dots) Steller sea lions counted on Walrus Island, Pribilof Islands. Data from National Marine fisheries Service, National Marine Mammal Laboratory, Seattle, Washington. From Hunt et al., 2002a.




Figure 27. Population trends in Black-legged Kittiwakes and red-legged Kittiwakes on the Pribilof Islands. From Dragoo et al., 2001.












Figure 28. Population trends of Common and Thick-billed Murres on the Pribilof Islands. From dragoo et al. 2001.













Figure 29 Changes in prey composition for northern Fur seals on the Pribilof Islands in the late 1980s and 1990s. Gm/Gm 'shelf squid', one or both of two species *Gonatus middendorffi* and/or *G. madokai*. Gb/Bm is 'slope squid' *Gonatopsis borealis* and/or *Berryteuthis magister*. SNP = St. Paul and SNG = St. George. From Sinclair et al., In Prep.









Figure 30. Changes in the age composition of walleye pollock in the diets of northern fur seals on the Pribilof Islands in the late 1980s and 1990s. Data from Sinclair et al., In Prep.






Figure 31. changes in the species composition of the diets of black-legged and red legged kittiwakes at the Pribilof Islands, 1975-1999. Data from Vlietstra et al. 2003.





Figure 32. Top: year class strength of walleye pollock; Bottom: biomass of pollock on the eastern Bering Sea shelf. (NPFMC, 2002)

# Oscillating Control Hypothesis




Figure 33: Schematic of the function of the Oscillating Control hypothesis. For an explanation, see text. From Hunt et al., 2002a).